

## Formulation and Evaluation of Oral Medicated Jellies of Psidium Guajava Linleaves

Bhosale Asmita Chandrakant<sup>1\*</sup>, Ms Darshana Varma<sup>2</sup>, Ms. Siddhi Chandak<sup>3</sup>,

*SMBT Institute of Diploma in Pharmacy<sup>1</sup>*

*Lecturer of SMBT institute of Diploma in Pharmacy<sup>2</sup>*

*SMBT College of Pharmacy<sup>3</sup>*

Submitted: 11-03-2024

Accepted: 21-03-2024

### ABSTRACT :

The guava leaves are obtained from the plant Psidium guajava l belong in to family Mrytaceae. Guava leaves have long been used effectively due to their antimicrobial, antibacterial and antiulcer properties .There are several type of jellies are available to treat ulcere like medicated jellies ,lubricant jellies, miscellaneous jellies . The aim for this research is prepared a oral medicated jellies with the help of guava leaves for the treatment of peptic ulcer .The oral route of administration is the most convenient and acceptable route for the patient compliance and acceptance .

**KEY POINTS** ; Medicated jellies , guava leaves ,peptic ulcer , anti ulcer property .

### I. INTRODUCTION:

Patients typically feel at ease using oral medication administration systems. Additionally, patient compliance may be improved by the oral drug delivery system's efficacy, safety, and effectiveness (1). The guava leaves in this study have a number of qualities, including antimicrobial, antibacterial, actions that are spasmolytic, analgesic, and antiulcer. Due to antiulcer property of guava leaves are use in preparation of oral medicated jellies. The main advantage of oral medicated jellies is it can administer easily and use the jellies anytime, anywhere as it is easy to handle and does not require water. (2)

### Types of jellies

#### 1. Medicated jellies:

Medicinal jelly Their spermicidal, local anesthetic, and antiseptic qualities are the main reasons they are applied to skin and mucous membranes. Enough water is available within these jellies. Jelly retains any residual protective layer and provides a localized cooling effect once the water has evaporated.(1)

#### 2. Lubricating jellies:

Medical devices used in diagnostic procedures, such as surgical gloves, cystoscopes, catheters, and rectal thermometers, are lubricated with these jelly-like substances. To be used as lubricants to introduce things into sterile bodily cavities, such as the pee bladder, layer jellies must be sterile.(3)

#### 3. Miscellaneous jelly

These are meant to be used for several purposes, such as patch testing and electrocardiography.(1)

#### Advantages of oral medicated jellies

1. The treatment is easily halted at any moment if necessary.
2. You don't need water to take your prescription.
3. Trustworthy, easy to use, and convenient administration
4. Boost the adherence of patients.
5. It is beneficial for those who have dysphagia.
6. Avoid first pass metabolism or increase bioavailability.
7. Pediatricians highly recommend it.
8. Well accepted by elderly and geriatric patients
9. It can be taken without water at any time or location.(2)

#### Objectives

1. Children voluntarily eat jelly candies because they enjoy the flavor and the fact that fruit juice and extract are used to flavor them, which leads to the development of jelly candies.
2. To develop a dosage form suitable for those suffering from dysphagia.
3. To improve the safety, efficacy, and bioavailability of the medication.
4. Creating dose forms that comply with patients is a better option than using conventional ones.(1)

#### Various componentof medicated jellies formulation

**I. Gelling Agent:** These hydrocolloids create matrices that resemble gel. It disintegrates in liquid phase and creates an internally cohesive structure that is weak. Illustrations of gelling agents:

a) Sodium Alginate: Brown algae's cell walls are the source of alginate. Alginates create thick gum when they bind with water. It is a component of several topical and oral medicinal preparations. It is typically employed in a variety of topical formulations, including pastes, creams, and gels, as a thickening and suspending agent.(4)

b) It is a heteropolysaccharide derived from terrestrial plant cell walls. It is used to treat diarrhea and constipation by making stools more viscous and volumetric.

Owing to its lower cost, it is utilized in a variety of drug delivery techniques, including mucoadhesive, controlled release, gastroretentive, and colon-specific drug administration systems. utilized as stabilizers in cosmetics as well.

c) Tragacanth • Tragacanth gum functions as an emulsifier and suspending agent in a range of medicinal formulations, including gels, creams, and emulsions.

Additionally utilized in foods and medications as an additive, thickener, and stabilizer. (5)

d) Gelatin • In pharmaceutical preparation, vitamin capsules, cosmetic technologies, and photographic emulsions, gelatin is typically utilized as a gelling agent.

utilized to administer medications suspended in biodegradable matrices in implanted delivery systems as well.(6)

## II. METHOD AND MATERIALS :

### Material:

Formulation of oral medicated jellies, the Guava leaves are received from local market in Nashik. Gelatin as gelling agent and other excipients like Sucrose, dextrose, citric acid as pH modifier and preservatives like methyl paraben and propyl paraben received from SMBT Institute of Diploma Pharmacy Nashik.

### Method :

1. Extraction of Guava leaves with the help of Soxhlet apparatus.

2. Preparation of oral medicated jellies With the help of Guava extract, gelling agent, preservatives, ph modifier and excipients. The oral medicated jellies prepared With heating and congealing method. Then evaluate the jellies with their evaluation test.(7)(8)



Fig 1- Psidium guajava



Fig no 2- Soxhlet Extraction



Fig no 3- Water Bath evaporation of Extract

**Formulation table:**

Ingredients	F1	F2	F3	F4	F5
Drug	0.04gm	0.04gm	0.04gm	0.04gm	0.04gm
Gelatin	2gm	2.50gm	3 gm	3.50gm	5gm
Citric acid	0.3gm	0.3gm	0.3gm	0.3gm	0.3gm
Methyl paraben	0.05gm	0.05gm	0.05gm	0.05gm	0.05gm
Propyl paraben	0.03gm	0.03gm	0.03gm	0.03gm	0.03gm
Dextrose	0.5gm	0.5gm	0.5gm	0.5gm	0.5gm
Sucrose	10gm	12.50gm	13.34gm	14gm	15gm
Water	5.73ml	6ml	5.48 ml	5.73 ml	6 ml

**Preparation of oral jellies**

1. weight the ingredient
- 2.in beaker mix sucrose with required amount of water and maintain temperature 80 degree
3. gelling agent add in that solution with contineou stirring

- 4.After mixing the gelling agent the ph modifier and preservatives must be added
- 5.then addrequired amount of main ingredient and mix before the gellies are going to set
- 6.finally transfer the formulation in the mold.(9)(10)

**Evaluation table**

**1. Physical evaluation**

Evaluation parameters	F1	F2	F3	F4	F5
Color	Light yellow	Light yellow	Light yellow	Light yellow	Yellow
Odour	Odourless	Odourless	Odourless	Odourless	Odourless
Taste	Aromatic	Aromatic	Aromatic	Aromatic	Aromatic
PH	3.3	3.4	3.3	3.2	3.4
Appearance	Transparent	Transparent	Acceptable	Acceptable	Acceptable
Consistency	Slightly liquid	Slightly liquid	Thick	Acceptable	Acceptable
Texture	Sticky	Sticky	Non sticky	Non sticky	Non sticky
Weight variation	9.47	9.49	9.78	9.93	9.52
Stickiness	Sticky	Sticky	Non sticky	Non sticky	Non sticky

2. Drug release

Time	F1	F2	F3	F4	F5
5	19.56	22.07	17.03	18.43	21.07
10	33.78	35.16	27.68	37.65	40.16
15	54.89	49.05	40.91	59.05	61.97
20	68.72	70.44	62.56	75.14	79.58
25	84.32	86.56	78.89	89.87	88.63
30	95.73	98.12	93.58	98.52	91.77

3. Sensory evaluation

Parameter	F1	F2	F3	F4	F5
Color	10	9	8.5	10	9
Taste	7.5	8.5	8	9	9
Flavour	8.5	8.5	8.5	9	10
Shape	8.5	9	7.5	9	8.5
consistency	8	8.5	8	10	9

Result:

The oral medicated jellies of guava leaves was prepared and evaluated. The guava leaves extract

was prepared in soxhlet apparatus and the all medicated jellies was prepared in heating and congealing methods.



III. CONCLUSION:

The development and assessment of oral medicated jelly as a gelling agent at higher, medium, and lower concentrations was the goal of the current investigation. The prepared jelly

batches (F1 through F5) were assessed for consistency, pH, and appearance. In this work, gelatin was successfully used to produce jelly that was filled with guava extract. The jellies are used as

an alternative to various solid dose forms for oral administration to juvenile and geriatric patients.

#### REFERENCE:

- [1]. Formulation And Evaluation Of Oral Soft Jelly Containing Glibenclamide, n.d.; Asmita 11, n.d.; Asmita 5, n.d.; Asmita 6, n.d.; Asmita 7, n.d.; Asmita 9, n.d.; Chitare et al., 2023; Gomaa & Ayoub, 2021; Ha et al., 2016; Honale et al., 2023; Kadam et al., 2020; Manali & Shah, 2016; Nayak et al., 2016; Prakash et al., 2014; Rathod et al., 2022)
- [2]. Chitare, S. S., Kailas Gosavi, T., & Patil, C. D. (2023). International Journal of Research Publication and Reviews FORMULATION AND EVALUATION OF HERBAL NANO GEL. International Journal of Research Publication and Reviews, 4(5), 316–325. www.ijrpr.com
- [3]. Gomaa, E., & Ayoub, M. M. (2021). Vardenafil oral jellies as a potential approach for management of pediatric irritable bowel syndrome. Saudi Pharmaceutical Journal, 29(9), 955–962. <https://doi.org/10.1016/j.jsps.2021.07.020>
- [4]. Ha, P. P., Nayak, K., Mishra, M. K., & Verma, G. (2016). I in Nd Do O a Am Me Er Ri Ic Ca an N J Jo Ou Ur Rn Na Al L O of F Formulation and Evaluation of Oral Soft Jelly Containing Glibenclamide. IAJPS 2016 J. P. Sci, 3(310), 1276–1282. <https://doi.org/10.5281/zenodo.167095>
- [5]. Honale, V. S., Muneshwar, S. D., & Sawale, A. V. (2023). Formulation and Evaluation of Oral Soft Jelly Containing Salbutamol Sulphate for the Treatment of Asthma. Journal of Drug Delivery and Therapeutics, 13(6), 118–124. <https://doi.org/10.22270/jddt.v13i6.6093>
- [6]. Kadam, V. S., Kendre, J., Shendarkar, G. R., & Kadam, S. S. (2020). Formulation and Evaluation of Medicated Oral Jelly of Trazadone Hydrochloride. International Journal of Pharmaceutical Sciences and Research, 11(12), 6251. [https://doi.org/10.13040/IJPSR.0975-8232.11\(12\).6251-59](https://doi.org/10.13040/IJPSR.0975-8232.11(12).6251-59)
- [7]. Manali, M. R., & Shah, D. P. (2016). Oral medicated jelly: a recent advancement in formulation. Pharma Science Monitor, 7(2), 13–20.
- [8]. Nayak, K., Mishra, M. K., & Verma, G. (2016). Formulation and Evaluation of Oral Soft Jelly Containing Glibenclamide. INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES, 3(10), 1276–1282. <https://doi.org/10.20959/wjpps20235-24699>
- [9]. Prakash, K., Satyanarayana, V. M., Nagiat, H. T., Fathi, A. H., Shanta, A. K., & Prameela, A. R. (2014). Formulation development and evaluation of novel oral jellies of carbamazepine using pectin, guar gum, and gellan gum. Asian Journal of Pharmaceutics, 8(4), 241–249. <https://doi.org/10.4103/0973-8398.143937>
- [10]. Rathod, M. P. I., Deshpande, P. V. V, Shaikh, S., Pathare, M. S. B., Nawghare, M. S. M., & S, M. S. (2022). Review on Formulation and Development of Oral Medicated Jellies. International Journal of Pharmaceutical Research and Applications, 7(4), 174–179. <https://doi.org/10.35629/7781-0704174179>