

A Review – Nanosponges for Antifungal Activity

Miss. Savita Dhamake, Miss.Dhanashri Vekhande, Miss. Dipali Thakare, Miss. Pranali Harad, Dr. Shoheb Shaikh

1. Research Scholar, Siddhi Institute of Pharmacy, Thane 421401

2. Research Scholar, Siddhi Institute of Pharmacy, Thane 421401

3. Research Scholar, Siddhi Institute of Pharmacy, Thane 421401

4. Research Scholar, Siddhi Institute of Pharmacy, Thane 421401

Date of Submission: 18-05-2024

Date of Acceptance: 28-05-2024

ABSTRACT:

The recent advance in nanotechnology has led to the development of targeted drug delivery system. However, targeting a molecule to a particular Site using a drug delivery system effectively requires a specialized drug delivery system. The discovery of Nano sponge has become a significant step in overcoming certain problems such as drug toxicity, poor bioavailability and release of drug in a predictable fashion as they can accommodate both hydrophilic and hydrophobic drug. Nano sponges exhibit a porous structure in nature which has the unique ability to entrap the drug moieties and offers a merit of desire release. Nano sponges are tiny sponges that can circulate in the body to reach the specific site and binds on the surface to Release the drug in a controlled and predictable manner. Nano sponges can be formulated by crosslinking of cyclohexatriene with carbonyl or decarboxylase (Cross linkers). Nano sponge's technology has been explored widely for the delivery of drugs for oral administration, topical Administration, and parental administration. Nano sponges can also serve as an effective carrier for enzyme, proteins, vaccine and antibodies. The Present review highlights the method of preparation, characterization and their potential application in drug delivery system.

Keywords: - Targeted drug delivery system, Nano sponges, Hydrophilic and Hydrophobic drug, β -Cyclohexatriene.

INTRODUCTION: -




Fig No.01 Structure of Nanosponges showing a cavity for drug loading

To get intended result, targeting drug delivery systems have been an ambition for a prolonged period. In the beginning, Nano sponge drug delivery system appeared only as a topical delivery system, but in the 21st century, Nano sponges can be administered by oral as well as intravenous (IV) route [1]. Nano sponge is a modern category of material and is made up of tiny particles with a narrow cavity of few nanometers. These narrow cavities can be filled with various types of substances. These tiny particles are having a capability due to which it is able to carry both hydrophilic and lipophilic drug substance and can increase the stability of poorly water-soluble drug substance or molecules [2]. The Nano sponges are a three-dimensional scaffold (backbone) or network of polyester that are capable of degrading naturally. These polyesters are mixed with across linker in solution to form Nanosponges. Here, the polyester is generally biodegradable, so it breaks down in the body moderately. Once the scaffold of Nano sponge's breaks down it releases the drug molecules which is loaded, in a derogatory fashion.

1.0.1 Advantages of Nano sponges:-

- i. Increase aqueous solubility of the poorly water-soluble drug.

- ii. Nano sponges can release the drug molecules in a predictable
- iii. Fashion.
- iv. Because of their tiny pore size (0.25 μm), bacteria cannot
- v. Penetrate the Nano sponges and they act like a self-sterilizer.
- vi. Nano sponge's drug delivery system are non-irritating, nonmutagenic and non-toxic.
- vii. Nano sponges help to remove the toxic and venom substance from the body.
- viii. Nano sponges drug delivery system minimize side effect.
- ix. Increase formulation stability and enhance the flexibility of the formulation.
- x. Reduce dosing frequency.
- xi. Better patient compliance.

1.0.2 Disadvantages of Nano sponges:-

- i. Nano sponges have the capacity of encapsulating small
- ii. Molecules, not suitable for larger molecules.
- iii. Dose dumping may occur at time.

• 1.2.0 Methods of Preparation: -

• 1.2.1 Solvent Method: -

Using solvent method, Nano sponges are prepared by mixing polar aprotic solvents like Dimethyl sulfoxide (DMSO), Dimethylformamide (DMF) with the polymer. A cross linker is then added to this mixture in the ratio of 1:4. The above reaction should be proceeded at temperature 10°C to reflux the temperature of the solvent for the time ranging from 1 to 48 h. Once the reaction has completed, the solution is cooled down at room temperature and then obtained a product is added to bi-distilled water. The product is recovered by filtering the product under vacuum and refining by soxhlet extraction with ethanol followed by drying.

• 1.2.2 Emulsion solvent diffusion Method: -

In this method, different proportion or amount of ethyl cellulose and polyvinyl alcohol are used to prepare Nano sponges. Two phases are used in this method—dispersed and continuous. The dispersed phase consists of ethyl cellulose and the drug, which is then dissolved in 20 ml of dichloromethane and some amount of polyvinyl alcohol (PVA) is added to 150 ml of the continuous Phase (aqueous). Then, the mixture is stirred at the speed of 1000 rpm for about 2 h. The product i.e. the Nano sponges are collected by filtration.

Finally, the product is dried in an oven at a temperature of 40°C .

• 1.3.0 Mechanism of drug release from Nano sponges: -

Since the Nano sponges have an open structure (in the surrounding of Nano sponges they do not have any continuous membrane), the active substance is added to the vehicle in an encapsulated form. The encapsulated active substance is able to move freely from the particles into the vehicle until the vehicle gets saturated and the equilibrium is obtained. As soon as the product is applied on to the skin, the vehicle containing the active ingredient gets unsaturated causing a disturbance in the equilibrium. Thus, the flow of active substances from Nano sponge particles into vehicles starts to epidermis until the vehicle is either absorbed or dried. Even after the retention of the Nano sponge particles on the surface of skin i.e. the stratum carenum, the release of active substance continues to skin for a long period of time.

• 1.4.0 Factors influencing in the formulation of Nano sponges:-

• 1.4.1 Nature of polymer: -

The polymer used in the preparation of Nano sponges can influence its formation and can also affect the pre-formulation. The size of the Cavity of a Nano sponge should be big enough to entrap a drug Molecule of a particular size into it for complexation.

• 1.04 Drug: -

- To be complex with Nano sponges, the drug molecules should have some specific characteristics as mentioned below:
- The molecular weight of the drug molecule should be in range ranging from 100-400 Daltons.
- Structure of drug molecule should not consist of more than 5 Condensed ring.
- The solubility of the drug in water should be $<10\text{ mg/ml}$.
- The melting point of the drug should be $<250^{\circ}\text{C}$.

• Temperature: -

Changes in the temperature can affect the complexation of drug or Nano sponges. Increasing the temperature generally decreases the extent of the stability constant of the drug or the Nano sponge complex which may be due to the reduction of interaction forces such as hydrophobic forces

and Van der Waal forces of drug/Nano sponges with an increase in the temperature.

- **Method of preparation: -**

The method of drug loading into the Nano sponges can cause a change in the complexation of drug and the Nano sponges. Although, the success of a method mainly depends on the nature or the characteristics of the drug and polymer; in some cases, freeze drying has also been known to affect the drug and Nano sponge complexation.

- **Characteristics of Nano sponges: -**

- Nano sponges are porous particles, used mainly to encapsulate the poorly soluble drugs.
- These Nano sponges have high aqueous solubility and are capable of carrying both lipophilic and
- Hydrophilic drugs.
- Nano sponge's formulations are stable over the pH range of 1 to 11 and temperature up to 300 °C
- Nano sponges are non-irritating and non-mutagenic, non-allergic and non-toxic and protect the drug from physiological degradation.
- Nano sponges can encapsulate various types of drug molecules by forming inclusion and non-emulsion complexes.

- **Particle size determination: -**

The particle size of Nano sponge is an important criterion in the optimization process of nanosponge. The particle size of the drug can affect the drug release as well as the solubility of the drug. Particle size can be determined by using the instrument, laser light diffractometric or Zeta sizer. Cumulative percentage drug release from Nano sponges of different particle size can be plotted against time to study the effect of particle size on drug release. Particle size larger than 30 µm can show gritty feeling and particle size range from 10 –25 µm can be preferred for topical drug delivery.

- **Solubility studies: -**Higuchi and Connors explained the method to study the inclusion complexation known as phase solubility method. This method used to explain the effect of Nanosponge on the solubility of the drug, which indicates the degree of complexation.

- **Microscopy studies: -**Scanning Electron Microscopy (SEM) and Transmission Electron Microscopy (TEM) can be used to study the

microscopic aspects of the drug and nanosponge formulation. SEM are used for the study morphology of the nanosponges. The difference in crystallization state of the raw materials used for the preparation of nanosponge and the final formulation seen under electron microscope indicates the formation of the inclusion complexes.

- **IR spectroscopy: -**

IR spectroscopy is used to estimate interaction between drug molecules and drug and nanosponge in the solid state. IR changes if there is complex formation between drug and nanosponge and if a small fraction of the drug molecule is encapsulated in complex less than 25 percent band and assigned to include part of another molecule which is marked by bands of the spectrum of nanosponges. IR has limited application to some drugs containing such as carbonyl or sulfonyl groups. IR study gives information of functional group containing drug.

- **Zeta Potential: -**

Zeta potential can be measured by using instrument zeta sizer, which is the measure of the surface charge of Nanosponges. Zeta potential is widely used for quantification of the magnitude of the electrical surface charge at the double layer. The zeta potential value more than 30mV indicates the good stability of formulation.

- **Loading efficiency: -**

The loading efficiency of a nanosponge particle can be determined by the estimation of drug loaded into the nanosponge using UV spectrophotometer and high-performance liquid chromatography method for the nanosponges. The loading efficiency of nanosponges can be calculated by using the following equation.

- **Applications of Nanosponges: -**
- **Solubility Enhancement: -**

$$LE = \frac{\text{Actual drug content in nanosponges}}{\text{Theoretical drug content}} \times 100$$

Nanosponges can improve the wettability and solubility of molecules with poor solubility in water. The drug can be molecularly dispersed in the nanosponge structure and then released as a molecule, avoiding the dissolution step. Therefore, the apparent solubility of the drug can be increased. Many formulations and bioavailability problems

can be solved by improving the solubility and dissolution of substances and nanosponges can greatly improve the solubility of drugs.

• **Nanosponges for Drug Delivery: -**

Nanosponges Are solid in nature and can be formulated into oral, Parenteral, topical or inhalation dosage forms.

For oral administration, the complex can be dispersed in a matrix suitable for the preparation of Capsules or tablets with excipients, diluents, Lubricants, and anti-caking agents. For parenteral Administration, the complex can simply be carried in sterile water, saline or other aqueous solutions. For topical administration, they can be effectively incorporated into topical hydrogels.

• **Topical Agents: -**

The nanosponge delivery system is a unique technology used to control the release of topical drugs that extend drug release and retention of drug form on the skin. Local anesthetics, antifungals, and antibiotics belong to the category of drugs that can be easily formulated into topical Nano sponges. When the active ingredient penetrates into the skin, a rash or more serious side effects may occur.

• **Nanosponges as a Carrier for Delivery of Gases: -**

Gas plays an important role in medicine, whether it is used for diagnostic or therapeutic

purposes. The Lack of an adequate supply of oxygen, called Hypoxia, is associated with various pathologies, from inflammation to cancer. In clinical practice, it is sometimes difficult to deliver oxygen in an appropriate form and dose. Cavalli et al. develop nanosponge formula as an oxygen delivery system for topical application, with the ability to store and release oxygen slowly over time.

• **Role of nanosponges for treatment of fungal: -**

• **Fungal Infection: -**

Fungal infections, or mycosis, are diseases caused by a fungus (yeast or mold). Fungal infections are most common on your skin or nails, but fungi (plural of fungus) can also cause infections in your mouth, throat, lungs, urinary tract and many other parts of your body.

Type of fungal infection: -

- i. **Superficial: -** Affect skin – mucous membrane. E.g. tinea versicolor dermatophytes: Fungi that affect keratin layer of skin, hair, nail. E.g. tinea pedis, ring worm infection Candidiasis: Yeast-like, oral thrush, vulvo-vaginitis, nail infections.
- ii. **Deep infections: -** Affect internal organs as: lung, heart, brain leading to pneumonia, endocarditis, and meningitis




Fig No:-02 [Types of fungal infection]

- **Overview of Fungal Skin Infections: -**
- **Physiology of normal skin**
- The skin is composed of three layers,
 - Epidermis (50–100 μm)
 - Dermis (1–2 mm)
 - Hypodermis (1–2 mm)




Fig.No.03 [Internal structure of skin]

Fungi usually make their homes in moist areas of the body where skin surfaces meet between the toes, in the genital area, and under the breasts. Common fungal skin infections are caused by yeasts (such as *Candida* or *Malassezia furfur*) or dermatophytes, such as *Epidermophyton*, *Microsporum*, and *Trichophyton*. Many such fungi live only in the topmost layer of the epidermis (stratum corneum) and do not penetrate deeper. Obese people are more likely to get these infections because they have excessive skin folds, especially if the skin within skin fold becomes irritated and broken down (intertrigo). People with diabetes tend to be more susceptible to fungal infections as well. Strangely, fungal infections on one part of the body can cause rashes on other parts of the body that are not infected. For example, a fungal infection on the foot may cause an itchy, bumpy rash on the fingers.

These eruptions (dermatophytids, or identity or id reactions) are allergic reactions to the fungus. They do not result from touching the infected area.



Fig no:-04 [Fungal infection on skin]

- **Symptoms:-**
 - i. Skin changes
 - ii. red and possibly cracking
 - iii. Peeling skin.
 - iv. Itching
- **Causes of fungal skin infection:-** Imbalance of bacteria is due to following reasons:-
 - Due to use of antibiotics
 - Hormone imbalance
 - Poor eating habits

- **Diagnosis:-**

Doctors may suspect a fungal infection when they see a red, irritated, or scaly rash in one of the commonly affected areas. They can usually confirm the diagnosis of a fungal skin infection by scraping off a small amount of skin and having it examined under a microscope or placed in a culture medium where the specific fungus can grow and be identified.

- **Treatment:-**

- Antifungal drugs
- Measures to prevent moisture
- Fungal infections are typically treated with antifungal drugs, usually with antifungal
- Drugs that are applied directly to the affected area (called topical drugs). Topical drugs
- May include creams, gels, lotions, solutions, or shampoos. Antifungal drugs may also be taken by mouth.
- In addition to drugs, people may use measures to keep the affected areas dry, such as
- Applying powders or wearing open-toed shoes.
- For some infections, doctors give corticosteroids to relieve inflammation and itching.

- **Introduction to Nanoemulgel:-**

- **Plant profile:-**

- **Pongamia Pinnata:-**

Pongamia pinnata (L.) Pierre belongs to the Fabaceae family, a medium-sized perennial tree commonly known in Hindi as Karanja and in English as Indian beech. *Pongamia pinnata* has been documented in a number of traditional medicinal systems for the cure of various human diseases and foods. It includes alkaloids, flavonoids, tannins, hormones, glycosides, karangin, glabrin, kanugin, and fixed oils, as well as other phytoconstituents. Historically, *P. pinnata* has been used as a folk medicinal plant, mostly in the Indian medicine systems of Ayurveda and Siddha. The anti-inflammatory, anti-nociceptive, antioxidant, anti-diarrhoeal, anti-fungal, anti-plasmodial, anti-ulcer, anti-hyperglycemic, anti-lipoxidative, anti-hyperammonemic and analgesic functions are available in plant extracts. The tree is known for its multipurpose advantages and as a potential biodiesel source.

- **Synonyms:-**

Pongamia pinnata is sometimes referred to by different synonyms, such as *Millettia pinnata* (L.) Panigrahi, *Derris indica* (Lam.), *Pongamia glabra*

Vent. Pongamia pinnata Merr. Bennett, *Millettia novo-guineensis* Kane. & Hat

- **Taxonomical Classification of Pongamia pinnata (L.)**

- ❖ Kingdom - Plantae
- ❖ Subkingdom - Tracheobionta
- ❖ Super division - Spermatophyta
- ❖ Division - Magnoliophyta
- ❖ Class - Magnoliopsida
- ❖ Subclass - Rosidae
- ❖ Order - Fabales
- ❖ Family - Fabaceae
- ❖ Genus - *Pongamia*
- ❖ Species – *Pinnata*
- ❖ Vernacular names – Different vernacular names of *P. pinnata* have been reported as follows
- ❖ Telugu-Kanuga
- ❖ Tamil-Punagai, dalkaramacha, pongam, punku
- ❖ Hindi-Karanja, pongam oil tree, paper, kanji
- ❖ Bengali-Karach
- ❖ Sanskrit-Naktamala
- ❖ Kannada-Honge, hulagilu
- ❖ Urdu-Karanja
- ❖ Malayam-Pungu, ungu, unu, avital
- ❖ Oriya-Koranj

- **Origin and geographical distribution :-**

It was naturally distributed in Asia, now this is found in Australia, Florida, Hawaii, India, Malaysia, Oceania, Philippines and Seychelles. It was commonly grown in coastal forest over India and near the streams and rivers.

- **Botanical description:-**

Pongamia pinnata is a fast-growing tree that ranges 40 feet in height & spreads, creating a wide-ranging, spreading canopy that casts moderate shade, according to Allen and Allen (1981).

Pongamia pinnata, a versatile resource of essential oil can be obtained through Soxhlet extraction method. The higher yield of seed extract can be achieved in the temperature range of 60-70°C. It has antifungal, antimicrobial properties which restrict the growth of microorganism as discussed in literature. The seeds of *Pongamia pinnata* contain recognised flavonols, have been reported. These biological features can be employed to generate new medication that can be used to replace those that are no longer effective.

- **Plant type: -**
 - A tree of medium height, evergreen, annual and deciduous.
 - Height: Thirty-five to forty feet
 - Texture: Medium
 - Growth rate: Fast
 - Chromosome number: 22
- **Phyto-chemistry:-**
 - **Contents of Leaves:** -Alkaloid, Carbohydrates, Flavonoids- Kaempferol, Quercetin, Rutin, Tannin, Saponin, Phytosterol.
 - **Contents of seeds:-**Karangin,Kaempferol ,Kanugin, Kankone ,Alkaloids dimethoxykanugin ,Glabrin ,Gamatay ,Glabrosaphonin ,Tannin, β -sitosterol ,Saponin, Quercetin ,Pongapin ,Pinnatin ,Pongamol ,Neoglabrin.
 - **Contents of seed oil: -**Karanjin (S18H22O4), Pongamol, Glabrachalcone, Kanjone, Pongapin
 - **Content of bark: -**Resin, Contains a bitter alkaloid, Sugar, Mucilage.



[A]



[B]



[C]



[D]

Fig No:-05 Pongamiapinnata: - [A] The whole plant [B] Leaves [C] Fruits [D] Seeds

- **NEEM:-**
Azadirachta indica commonly known as neem, is native of India and naturalized in most of tropical and subtropical countries are of great medicinal value and distributed widespread in the world. The chemical constituents contain many biologically active compounds that can be extracted from neem, including alkaloids, flavonoids, triterpenoids, phenolic compounds, carotenoids, steroids and ketones, biologically

most active compound is azadirachtin, it is actually a mixture of seven isomeric compounds labelled as azadirachtin A-G and azadirachtin E is more effective¹. Other compounds that have a biological activity are salannin, volatile oils, meliantriol and nimbin.

The antifungal activity of neem leaves against pathogenic fungi like-*Aspergillus flavus*, *Alternaria solani* and *Cladosporium*.5% aqueous leaf extract of neem was shown to cause inhibition in growth of six tested fungal pathogens (*Aspergillus fumigatus*, *Aspergillus Niger*, *Aspergillus terreus*, *candida albicans* and *microsporium gypserum*)

- ❖ **Taxonomical classification:** -The taxonomic classification of neem is as follows:
- ❖ Kingdom: Plantae,
- ❖ Order: Rurales,
- ❖ Suborder: Rutinae,
- ❖ Family: Meliaceae,

- ❖ Subfamily: Melioideae,
- ❖ Tribe: Melieae,
- ❖ Genus: *Azadirachta*,
- ❖ Species: *A.indica*

• **Active compounds of *Azadirachta Indica L. (Neem)***

Azadirachta indica L. (neem) shows therapeutic role in health management due to rich source of various types of ingredients. The most important active constituent is azadirachtin and the others are nimbolinin, nimbin, nimbidin, nimbidol, sodium nimbinate, gedunin, salannin, and quercetin. Leaves contain ingredients such as nimbin, nimbanene, 6-desacetylnimbinene, nimbandiol, nimbolide, ascorbic acid, n-hexacosanol and amino acid, 7-desacetyl-7-benzoylazadiradione, 7-desacetyl-7-benzoylgedunin, 17-hydroxyazadiradione, and nimbiol.



[A]



[B]



[C]



[D]

Fig No: -06 [Neem]

One of the popular Type of medicinal herb is peppermint (*Mentha piperita* L.), which is known as pudina in India. This is mainly grown in hilly areas and mainly in cold climatic regions. It is widely known to the world for its cooling and calming effects. It is mentioned in Ayurveda as it is used to maintain the balance of 'vata dosha' whereas is used to decrease the 'pitta' and 'kapha doshas'. Peppermint (*Mentha piperita* L.) is actually a crossed-hybrid mint of watermint (*M. aquatica* L.) and spearmint (*M. spicata* L.) It belongs to the family of 'Lamiaceae'. In Sanskrit it is known as 'rochani', which means taste perception improver]. There are found many significant uses of peppermint in various industrial purposes, as well as it is used to prepare tisane. Leaves of this plant are mainly used to made tisane. Peppermint (*Mentha piperita* L.) is among the renowned mainly single component herbal teas. India stands at the topmost producer and exporter country of mint oil among all over the world, which is produced from the leaves of it after extraction. Mint oil and its constituent's and derivatives can be used in different types of industrial purposes including food, pharmaceutical, perfumery and flavouring industry.

Therapeutic phytochemical compounds present in peppermint has a number of bioactive phytochemicals that have well defined and proven chemical structures and possess significant physiological roles. Some of these components have been listed along with their IUPAC name in the table no.

• **Chemical Components of Peppermint:-**

- Polyphenols Rosmaric acid, eriocitrin, cinnamic acid, caffeic acid etc.
- Flavonoids glycosides –Narirutin, luteolin-7-O-rutinoside,isorhoifolin, hesperidin etc
- Limonene 1-methyl-4-(1-methylethenyl)-cyclohexene
- Cineole – 1,3,3 -Trimethyl-2-oxabicyclo [2.2.2] octane
- Methone (2S,5R)-2-isopropyl-5-methylcyclohexanone
- Menthofuran 3,6-Dimethyl-4,5,6,7-tetrahydro-1-benzofuran
- Isomenthone (2R,5R)-5-methyl-2-propan-2-ylcyclohexan-1-one
- Carvone 2-Methyl-5-(prop-1-en-2-yl)cyclohex-2-en-1-one,
- Pulegone p-Menth-4(8)-en-3-one
- Menthyl -acetate

- Acetic acid [(1R, 2S, 5R) -2-isopropyl-5-methylcyclohexyl] ester.

• **Role of peppermint:-**

Pharmacological effect Anti-fungal activity Pattnaik et al. (1996) found almost 11 to 12 types of fungus (including *Candida albicans*, *Trichophyton mentagrophytes*, *Aspergillus fumigatus* etc.) against which peppermint oil has shown significant inhibitory activity According to Nilo et al. (2017) the extract of peppermint leaves has fungicidal effect against the aggrandizement of some fungus including *Fusarium moniliforme*, *Aspergillus niger* and *Aspergillus fumigatus* According to Hussain et al. (2010), peppermint oil was found to show an inhibitory effect on *Aspergillus niger*, *Aspergillus flavus*, *Rhizopus solani* and *Fusarium solani* . Sandasi et al. (2011) suggested that the peppermint oil was found to inhibit the growth of *Candida albicans* and *Candida albicans* there are many relevant significant literature about peppermint and its fungicidal effect against different types of fungus and these have proved mechanism of action. Peppermint has been suggested for a number of antifungal treatments in human beings.

• **Gel:-**

The topical delivery of drugs is an attractive method for local and systemic treatments and is commonly used in the treatment of inflammatory conditions like dermatological diseases Topical application has many advantages over the conventional dosage forms, especially to avoid some serious systemic adverse effects A wide variety of pharmaceutical dosage forms can be used in the delivery system for topical drugs. The most used ones are gels, creams, and ointments, followed by sprays and liquid preparations the topical delivery with gels can increase the resistance time of the drug on the skin and improve the delivery and release of the substance by increasing the residence time at the injection site.

Gel formulations are used to deliver the drug topically because of easy application, increase contact time and minimum side effects as compare to other topical preparation and oral administration.

• **Preparation of gel base:-**

Carbopol 934 was dissolved slowly with stirring in 60 mL of demineralized water for 1 h to avoid agglomeration Then disodium edetate and triethanolamine were dissolved in 10 mL of demineralized water separately and stirred for 10

min. Mixed 4.83 mL of propylene glycol in 12 mL of demineralized water with stirring for 10 min. Disodium edetate and triethanolamine solution were added to carbopol solution and the pH was then adjusted to 7.4 by stirring the solution for 10 min. Then propylene glycol solution was added with stirring for 10 min until a clear consistent gel base was obtained.

• **Emulgel:-**

Emulgel is a combination of emulsion and gel, which is a new approach for topical delivery of

drug. It has a double control release like emulsion and gel. Gel is new class of formulation, it release the drug faster in comparison ointment, cream and lotion. Incorporation of drug in emulgel formulation is suitable to treat skin disorders. Emulgel -Emulgel having advantages of both gels and emulsion act as a controlled drug delivery system for typically applied Drug They are emulsion of either oil in water type or water in oil type which are gelled by mixing with a gelling agent.




Fig no:-07 [structure of Emulgel]

Emulgel is prepared both in oil- in- water and water- in-oil type emulsion mixed with gel. Oil- in- water type is used for lipophilic drugs and water- in- oil type is used for hydrophobic drugs' delivery. Emulgel is prepared both in oil- in- water and water- in-oil type emulsion mixed with gel. Oil- in- water type is used for lipophilic drugs and water- in- oil type is used for hydrophobic drugs' delivery.

• **ADVANTAGES:-**

- i. Incorporation of hydrophobic drugs
- ii. Better loading capacity
- iii. Better stability
- iv. Controlled release
- v. No intensive sonication
- vi. Avoiding first pass metabolism
- vii. Avoiding gastrointestinal incompatibility
- viii. More selective for a specific site
- ix. Improved patient compliance
- x. Convenient and easy to apply

- xi. Avoidance of the Systemic adverse effect the Systemic adverse effect of drug first pass metabolism in the body
- xii. Systemic circulation is minimize or prevented
- xiii. Improve patient compliance and acceptability
- xiv. Suitable for self-medication
- xv. Provide Target drug delivery on the body
- xvi. Ability to easily terminate medication when needed
- xvii. Can easily pass through skin having dual behaviour i.e hydrophobic as well as hydrophilic
- xviii. They are convenient to apply on hairy skin due to absence of greasiness and lack of residues upon application
- xix. Better stability and release of drug
- xx. Better loading capacity
- xxi. Production possibility and low preparation cost
- xxii. No intensive sonication needed

- xxiii. Emulgel can be used to prolong the effect of drug having shorter
- xxiv. Substrate for Oral absorption
- xxv. PH associated deactivation.

DISADVANTAGES:-

- i. Skin irritation on contact dermatitis
- ii. The possibility of allergenic reactions
- iii. The poor permeability of some drugs through the skin
- iv. Drugs of large particle size are not easy to absorb through the skin
- v. The occurrence of the bubble during formulation of emulgel

FORMULATION OF EMULGEL:-

For the preparation of emulgel some constituents are used including drug, which are:

❖ Vehicle:-

Vehicle should follow the ideal characters given in the Pharmacopeias

❖ Aqueous material:-

The aqueous phases used are water, alcohol, etc.

❖ Oil :-

Oils are used for preparation of emulsion. Mineral oils and paraffin are used either alone or in combination.

❖ Emulsifiers:-

Emulsifiers used for preparation of emulsion. Some examples are span 80, tween 80, stearic acid, sodium stearate.

❖ Gelling agents:-

Gelling agents are used for prepare gels, which enhance consistency of preparation.

❖ Penetration enhancers:-

Penetration enhancers help to absorb drug to the skin

❖ pH adjusting agent:-

IDEAL PROPERTIES OF ADDITIVES:-

- They should be nontoxic.
- They should be easily available.
- They should be cheap.
- They do not be contraindicated.
- They should be chemically and physically be stable.





Fig no - 08 [Preparation of emulgel]

PREPARATION OF EMLGEL:-

Emulgels are prepared by incorporating gel and emulsion. The emulsion and gel are prepared separately and mixed together. For preparing emulsion, aqueous phase and oil phase are taken separately and mixed together. Then the gel is prepared by using gelling agent. After preparing gel and emulsion, they are mixed with gentle stirring. The chemicals used as oil phase are castor oil, clove oil, liquid paraffin, etc. Water and alcohol are used as aqueous phase.

REFERENCE:-

- [1]. HIMANGSHU BHOWMIK, D. NAGASAMY VENKATESH, ANUTTAM KUILA, KAMMARI HARISH KUMAR Department of Pharmaceutics, JSS College of Pharmacy, Rockland's, Ooty 643001, JSS Academy of Higher Education and Research Received: 29 Jan 2018, Revised and Accepted: 24 May 2018.
- [2]. NANOSPONGES AS NOVEL CARRIER FOR TOPICAL DELIVERY OF LULICONAZOLE -ANTIFUNGAL DRUG. S. Wakure *, M. A. Salunke, P. T. Mane, S. R. Awale, R. D. Shinde and S. S. Eklinge Vilasrao Deshmukh Foundation, Group of Institutions, VDF School of Pharmacy, Latur - 413531, Maharashtra, India. Received on 15 July 2021; received in revised form, 25 August 2021; accepted, 28 August 2021; published 01 October 2021
- [3]. https://www.ncbi.nlm.nih.gov/core/lw/2.0/html/tileshop_pmc/tileshop_pmc_inline.html?title=Click%20on%20image%20to%20zoom&p=PMC3&id=9325902_gr1.jpg
- [4]. A Review on Pongamia pinnata (L.): Traditional Uses, Phytochemistry and Pharmacological Properties Akshay G. Fugare *1, Rajkumar V. Shete 2, Vishal S. Adak 3, Krishna Murthy G. 4 Received 27 Nov 2020;
- [5]. Antifungal Activity of leaf extract of Neem (Azadirachta Indica Linn) D.K. Shrivastava and Kshma Swarnkar* Govt. E. Raghavendra Rao Postgraduate Science College Bilaspur (Chhattisgarh) - 495001, India *Corresponding author ISSN: 2319-7706 Volume 3 Number 5 (2014) pp. 305-308.
- [6]. Journal of Survey in Fisheries Sciences 10(2S) 4063-4067 20234063 Evaluation of antifungal activity of .Department of Botany, Shahir Annabhau Sathe Mahavidyalaya Mukhed Dist. Nanded M.S., vijay.gorgile@gmail.com.
- [7]. World Journal of Pharmaceutical Research SJIF Impact Factor 8.084 Volume 11, Issue 1, 1375-1390. Research Article ISSN 2277- 7105
- [8]. DEVELOPMENT AND EVALUATION OF ITRACONAZOLE NANOSPONGES AS A FUNGI EVACUATOR -AN OPTIMISTIC DRUG DELIVERY SYSTEM Limce Thampi1, K. R. Akshay2, Gayathri Gopi2, Nazrin K. I.2, Ponnu Jacob2, Gini EJ.
- [9]. A phyto pharmacological review on a versatile medicinal plant: Pongamia pinnata (L.) pierre Sowjanya Pulipati, P Srinivasa Babu, D Neelima Lakshmi, N Navyasri, Y Harshini, J Vyshnavi and M Prasanth.
- [10]. A phyto pharmacological review on a versatile medicinal plant: Pongamia pinnata (L.) pierre Sowjanya Pulipati, P Srinivasa Babu, D Neelima Lakshmi, N Navyasri, Y Harshini, J Vyshnavi and M Prasant journal of Pharmacognosy and Phytochemistry 2018; 7(4): 459-463
- [11]. Antifungal effect of aqueous and ethanolic extracts of neem leaves, stem bark and seeds on fungal rot diseases of yam and cocoyam Chukwuma S. Ezeonu1*, Chinedu Imo1, Dawn I. Agwaranze2, Assumpta Iruka1 and Abraham Joseph.
- [12]. Bio-activity of oils of Trigonella foenum-graecum and Pongamia pinnata Pritee Wagh 1, Mahendra Rai*, S K Deshmukh 2 Marta Cristina Teixeira Durate 3 Department of Biotechnology SGB Amravati University Amravati-444602, Maharashtra, India. Department of Natural Products, Nicholas Piramal Research Centre, 1A, 1B&1C, Nirlon Complex, Off Western Express Highway, Next to Food Bazar, Goregaon (East), Mumbai 400 063. India. Biológicas E Agrícolas – Universidade Estadual de Campinas, Campinas - São Paulo, Brasil, P.O. Box 6171, CEP: 13081-970. Accepted 20 October, 2006.
- [13]. FABRICATION AND CHARACTERIZATION OF MICROSPONGE IMPREGNATED GEL

- FOR TOPICAL DELIVERY OF LULICONAZOLE Article in the International journal of analytical and experimental modal analysis · March 2022.
- [14]. A Review on Pongamia pinnata (L.): Traditional Uses, Phytochemistry and Pharmacological Properties Akshay G. Fugare *1, Rajkumar V. Shete 2, Vishal S. Adak 3, Krishna Murthy G. 4 Journal of Drug Delivery & Therapeutics. 2021; 11(1-s):207-211
- [15]. FUNGITOXIC PROPERTIES OF Pongamia pinnata (L) PIERRE EXTRACTS AGAINST PATHOGENIC FUNGI Digamber R. More and Mirza M. Vaseem Baig2* 1 Department of Botany, L.B.S.College, Dharmabad. Nanded. 431809 Maharashtra, India 2 Department of Botany, Yeshwant Mahavidyalaya, Nanded. 431602 Maharashtra, India *Corresponding author: Email: mmvbaig@gmail.com Tel: +91-9422170641; Fax +91- 2462-253726 [Received-11/10/2013, Accepted-26/12/2013].
- [16]. Formulation and evaluation of cyclodextrin-based nanosponges of griseofulvin as pediatric oral liquid dosage form for enhancing bioavailability and masking bitter taste Samia M. Omar a,b, Fares Ibrahim , Aliaa Ismail Received 21 October 2019; Accepted 29 January 2020 ; Available online 3 February 2020 .
- [17]. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9325902/>.
- [18]. Nanosponges: A Novel Targeted Drug Delivery for Cancer Treatment Tukaram S. Patil*, Nishigandha A. Nalawade, Vidya K. Kakade, Sumedha N. Kale.
- [19]. A Research Article on Nanogel as Topical Promising Drug Delivery for Diclofenac sodium Swati Talele1, Preetam Nikam1, Braja Ghosh1, Chaitali Deore2, Ashwini Jaybhav2, Anil Jadhav2
1 Department of Pharmaceutics, Sandip Institute of Pharmaceutical Sciences, Nashik, INDIA.
2 Department of Quality Assurance, Sandip Institute of Pharmaceutical Sciences, Nashik, INDIA.
- [20]. Formulation And Evaluation Of Emulgel Of An Antifungal Drug For Topical Drug Delivery Dinesh Kumar1*, Rajni2, Rahul Saini3, Suman Rani4, Raman Kumari5
1*Atam Institute of Pharmacy, Hisar, Haryana 2 Starex University, Gurgaon, Haryana 3Choudhary Bansilal university, Bhiwani 4Atam Institute of Pharmacy, Hisar, Haryana 5Atam Institute of Pharmacy, Hisar, Haryan *Corresponding Author: Dinesh Kumar.
- [21]. International Journal of Advanced Multidisciplinary Research ISSN: 2393-8870www.ijarm.com (A Peer Reviewed, Referred, Indexed and Open Access Journal)DOI: 10.22192/ijarmr Volume 10, Issue 4 -2023.
- [22]. Formulation of Entandrophragma utile into an Herbal Emulgel for the Management of InflammationAdeola Tawakalitu Kola-Mustapha 1,2,* , Haneefat Folashade Ibraheem 2, Suleiman Taiwo 2, Ismail O. Ishola 3 Sukurat Olasumbo Usman 4 and Yusuf Oluwagbenga Ghazali 2
- [23]. Formulation and Evaluation of Herbal Emulgel from CocciniaGrandis Linn Fruit Extract for Fungal Dermatological Infection.
- [24]. International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR) | February 2019 | Volume 9 | Issue 1 | Page 92-97 Sreevidya V.S, An Overview on Emulgel
ISSN (Online) 2249-6084 (Print) 2250-1029 www.eijppr.com An Overview on Emulgel
Sreevidya V.S Sreekrishna College of Pharmacy And Research Centre Parassala, Thiruvananthapuram, Pin: 695502, Kerala, India.
- [25]. Brazilian Journal of Pharmaceutical Sciences vol. 52, n. 3, Jul. /sep., 2016<http://dx.doi.org/10.1590/S1984-82502016000300015>Formulation and evaluation of topical herbal gel for the treatment of arthritis in animal model Rajasekaran Aiyalu*, Arulkumaran Govindarjan, Arivukkarasu Ramasamy KMCH College of Pharmacy, Coimbatore, Tamilnadu, India.
- [26]. World Journal of Pharmaceutical Research SJIF Impact Factor 8.084 Volume 11, Issue 2, 1657-1665. Research Article ISSN 2277– 7105
- [27]. FORMULATION AND EVALUATION OF HERBAL EMULGEL FOR

- TREATMENT OF FUNGAL INFECTIONS
Bhavyashree T.*, Shreya H. Alva, Shubhashree A. S., Spoorthi C., Sushmitha Rai, Shefali Bhute and A. R. Shabaraya Department of Pharmaceutics, Srinivas College of Pharmacy, Valachil, Mangalore, Karnataka, India.
- [28]. Wakure et al., UPSR 2022, Vol 12(10) 5570-5583 E-ISSN: 0975-8232, P-ISSN: 2320 5148
IJPSR (2021), Volume 12, Issue 10
- INTERNATIONAL JOURNAL OF PHARMACEUTICAL SCIENCES RESEARCH
Received on 15 July 2021 emed so remed fann 25 August 2121 accepted, 28 Augat 2021 published 01 October 2021
- NANOSPONGES AS NOVEL CARRIER FOR TOPICAL DELIVERY OF LULICONAZOLE-AN ANTIFUNGAL DRUG B. 5. Wakure, M. A. Salunke, PT Mane, 5 R Awale, RD Shinde and SS Eklinge
- [29]. The Scientific World Joutsal Volume 2016. Article ID 7394645, 4 pages
Research Article Development and Evaluation of Stability of a Gel Formulation Containing the Monoterpene Borneol Milla Gabriela Belarmino Dantas, Silvio Alan Gonçalves Bomfim Reis, Camila Mahara Dias Damasceno, Larissa Araújo Rolim, Pedro José Rolim-Neto, Ferdinando Oliveira Carvalho, Lucindo José Quintans-Junior, and Jackson Roberto Guedes da Silva Almeida.
- [30]. WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
Volume 11, Istur 1, 1876-1390)
Research Article ISSN 12777105
DEVELOPMENT AND EVALUATION OF ITRACONAZOLE NANOSPONGES AS A FUNGI EVACUATOR-AN OPTIMISTIC DRUG DELIVERY SYSTEM
Lince Thampi, K. R. Alahay, Gayathri Gopi, Nazrin K. 13, Ponnu Jacob, Gini Et'Departinent of Pharmaceutics, Chemists College of Pharmaceutical Sciences and Research, Vankoli PO. Puthencruz, Emkulan Kerala, lodia-682308 Chemists College of Pharmaceutical Sciences and Research,
- Vankoli PO. Puthencriz Ernakulam Kerala, India-682308
Department of Pharmacognosy, the Dale View College of Pharmacy & Research Centre, Punalal PO, Thiruvananthapuram, Kerala, India-695575
- [31]. WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
Volume 11, Istur 1, 1876-1390)Research ArticleISSN 12777105
DEVELOPMENT AND EVALUATION OF ITRACONAZOLE NANOSPONGES AS A FUNGI EVACUATOR-AN OPTIMISTIC DRUG DELIVERY SYSTEM
Lince Thampi, K. R. Alahay, Gayathri Gopi, Nazrin K. 13, Ponnu Jacob, Gini Et'Departinent of Pharmaceutics, Chemists College of Pharmaceutical Sciences and Research, Vankoli PO. Puthencruz, Emkulan Kerala, lodia-682308 Chemists College of Pharmaceutical Sciences and Research, Vankoli PO. Puthencriz Ernakulam Kerala, India-682308 Department of Pharmacognosy, The Dale View College Of Pharmacy & Research Centre, Punalal PO, Thiruvananthapuram, Kerala, India-695575.
- [32]. Original article Formulation and evaluation of cyclodextrin-based nanospheres of griseofulvin as pediatric oral liquid dosage form for enhancing bioavailability and masking bitter taste Samia M. Omar, Fares Ibrahim", Aliaa Ismail.
- [33]. Journal is Pharmacogeny and Phytochemistry 2018-4-40
Journal of Pharmacognosy and Phytochemistry Available online at www.phytojournal.com
21-15 2RIE A phyto pharmacological review on a versatile medicinal plant: Pongamia pinnata (L.) pierre Sanjanya Fuligati Sowjanya Pulipati, P Srinivasa Babu, D Neelima Lakshmi, N Navyasri, Y Harshini, J Vyshnavi and M Prasanth.
- [34]. Plant Archives Volume 21, No 1, 2021 pp. 515-525 ISSN: 2381-606) (online), ISSN: 0972-5210
Plant Archives Journal home page: www.plantarchives.org DOI <http://deler/1031470/PLANTARCHIVES201211072>

- FORMULATE AND EVALUATE GEL CONTAINING NANOSTRUCTURED LIPIDS OF PONGAMIA PINNATA Edakulathur Richou", Wanl Sucha Bhagyashree S. Mundhe, Saurabh Arun Rayate, Aishwarya Sahebrao, Talale Swati, and fadlaw Anll Deartment of Pluracetic Sandip Inte of Pharmaceutical Sciences, Nashikarana (22213, India Eñ npy 2018@gmail.com
Date of Racewing-01 11-2020, Date of Acceptance 31-01-2021
- [35]. Alncan Journal of Biotechnology Available online Vol. 6 (13), pp. 1500-1500, 4 July 2007 at hopu/www.academicjournals.org/AB ISSN 1684-5315 © 2007 Academic Joumata Full Length Research Paper Bio-activity of oils of Trigonella foenum-graecum and Pongamia pinnata Pritee Wagh, Mahendra Rai, SK Deshmukh and Marta Cristina Teixeira Durate
- [36]. Wakare ET ol, UPSR 2022, Vol 12(10): 5570-5583 E-ISSN 0975-8232, P-ISSN. 2320-5148
LJPSR (2021), Volume 12, Lastue 10
INTERNATIONAL JOURNAL
PHARMACGUTICAL SCIENCES
RESEARCH
Receved on 15 July 2021 din revised forum 25 August 2021 accepted 25 August 2021 published Oober 221
NANOSPONGES AS NOVEL CARRIER FOR TOPICAL DELIVERY OF LULICONAZOLE-AN ANTIFUNGAL DRUG IBS Wakre, MA Salunke, PT Mane, S R. Awale, R. D. Shunde and S. S. Ellinge
- [37]. FABRICATION AND CHARACTERIZATION OF MICROSPONGE IMPREGNATED GEL FOR TOPICAL DELIVERY OF LULICONAZOLE Article in the International journal of analytical and experimental modal analysis March 2022.
- [38]. Formulation and evaluation of herbal gel Article in Indian Journal of Natural Products and Resources December 2012 CITATIONS 3 authors, including: Vijay Gulkari J.L. Chaturvedi College of Pharmacy
- [39]. ANTIFUNGAL ACTIVITIES OF SEED OIL OF NEEM (Azadirachta indica A. Juss.) Article March 2014
3 authors: Adiyata Olutem Acepaja African Methodist Episcopal University 44 PUBLICATIONS 297 CITATIONSTunde Joseph GUNKLINLE Ladoke Akintola University of Technology
- [40]. The Pharma Innovation ESKY (E) 2277-269 Sunandini Wood Sornce and Technology, Voters Howeank Iostitate (Deed to Be Univermay Dehradum. Exurakhand, India Antifungal efficiency of Karanj seed extract.
- [41]. Evaluation of antifungal activity of Pongamia Pinnata L on Saprolegnia parasitica isolated from cultured and wild Fishes Vijay Tulshiram Gorgile Department of Botany, Shahir Annabhau Sathe Mahavidyalaya Mukhed Dist. Nanded MS. vijay gorgile@gmail.com
- [42]. International Journal of Current Microbiology and Applied Sciences ISSN: 2319-7706 Volume 3 Number 5 (2014) pp. 305-308 <http://www.ljcmas.com> Original Research Article Antifungal Activity of leaf extract of Neem (Azadirachta Indica Linn) D.K.Shrivastava and Kshma Swarnkar Govt. E. Raghavendra Rao Postgraduate Science College Bilaspur (Chhattisgarh) - 495001, India
- [43]. Journal of Chemical and Pharmaceutical Research and Prama J. Chem. Pharm. Res., 2010, 2(3):182-186ISSN No: 0975-7384 CODEN (USA): JCPRCS Formulation and evaluation of anti-bacterial and anti-fungal activity of a herbal ointment containing Aloe-vera, Azadirachta indica and Curcuma-long Abhijeet Pandey, Jui V. Jagtap, Aditi A. Patil, Richa N. Joshi and B. S. KuchekarDepartment of Pharmaceutics, MAEER's Maharashtra Institute of Pharmacy, Kothrud, Pune, MS, India.
- [44]. Antifungal effect of aqueous and ethanolic extracts of neem leaves, stem bark and seeds on fungal rot diseases of yam and cocoyam Chokwuma S Leanu" Chineda Ima Dawn! Agwarance Assumpta luka and Abraham Joseph.
- [45]. International Journal of Advanced Biotechnology and Research ISSN 0976-2612, Online ISSN 2278-599X, Vol 4,



Issue 4, 2013, pp 560-567

<http://www.bipublication.com>

FUNGITOXIC PROPERTIES OF
Pongamia pinnata (L) PIERRE
EXTRACTS AGAINST PATHOGENIC
FUNGI Digamber R. More' and Mirza M.
Vaseem Baig"Department of Botany,
L.B.S.College, Dharmabad Nanded.
431809 Maharashtra, India Department of
Botany, Yeshwant Mahavidyalaya,
Nanded. 431602 Maharashtra, India