International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 1125-1132 www.ijprajournal.com
ISSN: 2456-4494

Current Aspects and Future Potential of Transdermal Drug Delivery System

Samarth S. Kale Yogesh B. Raut Sanjay K. Bais

Date of Submission: 05-12-2025 Date of Acceptance: 12-12-2025

ABSTRACT:
The best way to donate pharmaceutical is one that is simple for patients to utilize, like taking pills by mouth, but it ought to too work as well as giving medication specifically into the blood, like through a shot. As of late, putting medication through the skin, called transdermal sedate conveyance (TDD), has gotten to be more prevalent. This strategy is not obtrusive, so it's less demanding for individuals who are more touchy, like children and more seasoned grown-ups, to utilize. It moreover makes a difference dodge a few issues that happen when taking medication by mouth, like the body not retaining it well or breaking it down as well much. To make medication pass through the skin superior, diverse strategies have been created. These incorporate inactive ways like utilizing little bubbles, minor particles, or extraordinary blends, and chemical ways like utilizing warm or weight. Researchers are too considering blended strategies that utilize both physical and chemical methods to offer assistance medication go through the skin more successfully. This paper surveys the current ways individuals are moving forward sedate conveyance through the skin, along with the great and not-so-good parts of these strategies.

Keywords: Barrier for the skin ;Topical ; drug Release ;Viability ;Nanoneedles ;microparticles ; Biological Nuclei ; Electrophoresis.

I. INTRODUCTION

Transdermal medicate conveyance a parcel of consideration from 20 years ago its numerous benefits comparatively to sedate conveyance strategies, such as its ease of utilize, pre-established measurements, ease of taking care of, persistent self-administration, and generally loose capacity conditions. In any case, the verbal course of medicate conveyance has a number of drawbacks when it comes to helpful compounds with restricted solvency, penetrability, and corruption. Since most solutions taken orally go through first-pass digestion system and confront various organic and physical deterrents, their verbal bioavailability changes broadly verbal medicate conveyance is the most effective strategy, particularly when persistent compliance cannot be ensured (e.g., comatose patients). In any case, specific devices, sterilizing strategies, and qualified staff are required for the generation and organization of parenteral details. Also, the request for a cheap, non-invasive immunization procedure, particularly in immature countries, has provoked seriously investigate into the creation of Consider substitute strategies that do not require needles, such transdermal sedate organization (TDD) [1].

TDD is a medicate conveyance strategy that is secure, well-tolerated, and is competent of combining the dose Verbal measurement shapes are related with exactness and ease of organization, though parenteral medicine conveyance is related metabolic free exchange for helpful specialist towards plasmatic state. Be that as it may, due

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 1125-1132 www.ijprajournal.com
ISSN: 2456-4494

to the moo skin porousness of a few drugs, its utility in modern clinical hone is restricted. The require to see dermis as a physical obstruction driven for the improvement of various strategies to move forward the transdermal entrance of drugs. This article points to donate a diagram of later progressions in TDD strategies and examines the focal points and impediments of the different dynamic (physical), inactive (chemical), and half breed strategies as of now being examined for making strides medicate penetrability over the skin[2].

While transdermal patches can be securely and effectively managed to children and the elderly TDD offers numerous points of interest over verbal sedate conveyance, counting dodging firstpass digestion system, protecting touchy medicines by unforgiving situation of GIT, Empowering supported discharge of drugs, in this way keeping up a more uniform plasma concentration [3].

Additionally, transdermal patches have been appeared to progress adherence in elderly polypharmacy patients who regularly complain of destitute compliance and trouble gulping since of the overwhelming every day pill burden . these skin steps - 1 epidermis, 2 dermis ,3 hypodermis as shown in figure1

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 1125-1132 www.ijprajournal.com
ISSN: 2456-4494



Figure 1. Overview of skin layer

In any case, in spite of all of its benefits, TDD has downsides. Incomprehensibly, the skin itself is the greatest impediment to TDD; the medicine interior a TDD framework must to begin with pass through the layers of skin in arrange to be retained into the systemic circulation [4].

The essential deterrent to TDD is the stratified physiological structure of the skin, and as it were a little number of medications—those with specific physicochemical characteristics—can enter the. skin and enter the plasma .A number of strategies and definition approaches that can offer assistance get past the skin obstruction have developed as a result of later advancements in transdermal medicine conveyance. These strategies are secured in the areas that follow[5].

II. PRESENTLY DECLARED TRANSDERMAL MEDICATIONS

The dissemination of numerous drugs has been altogether affected by the TDD advertise, basically in the zones of torment treatment hormonal applications clutters of the central apprehensive system cardiovascular illnesses and other applications, such stopping smoking . It is anticipated that the worldwide TDD advertise would be sizable. This industry is growing since to variables such the recurrence of inveterate ailments and headways in TDD procedures. The Joined together States affirmed the to begin with transdermal fix for systemic dispersion in 1979. 3-day fix managed nausea and motion sickness to remedy movement affliction. Buprenorphine is the most later fix endorsed for serious torment. endorsed by the FDA for the treatment of determined torment that doesn't move forward with other drugs [6].

Antiallergic begin with anti-histamine transdermal fix planned to treat unfavorably susceptible rhinitis, was affirmed for deal in Japan in 2018 . It has a long-lasting affect that endures up to 24 hours after ingestion and an effect that reduces allergy symptoms based on the amount taken[7].

In 2007, The FDA endorsed the to begin with Rotigotine-containing Parkinson's fix (NeuproTM,

UCB, Brussels, Belgium), a once-daily fix accessible in four dosing qualities: 2,4,6 mg[17]. To treat Brain infection, rivastigmine as

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 1125-1132 www.ijprajournal.com
ISSN: 2456-4494

of now legal to utilize as beneath the skin (ExelonTM, Novartis) . This fix disposes of the gastrointestinal (GI) side impacts connected to verbal rivastigmine [8].

An FDA-approved transdermal prophylactic fix containing norelgestromin and ethinyl estradiol is called Ortho EvraTM. For three weeks (21 days), the fix is connected once a week; the cycle incorporates one week without the fix [7,16]. ApleekTM is a transdermal prophylactic fix that contains 2.10 mg of gestodene and 550 mcg of ethinyl-estradiol as dynamic components. It is utilized once a week for three weeks, after at that point a patch-free interim of seven days [9].

III. STRATEGIES FOR PROGRESSING SKIN PERMEABILIZATION

TDD is restricted to a few medicines with specific physicochemical characteristics, in spite of its numerous points of interest. Where water loving and Karger molecular drugs like peptides, are regularly prevented by stratum corneum (SC) obstruction, a medicate candidate for transdermal conveyance to in a perfect world having a atomic weighing 500Da further direct to pass openly through the skin[10].

The essential defense from surrounding is subcutaneous of a skin layer that is 5–20 μm thick. Since ten to fifteen layers of framework, and close intersections that make up its structure (Figure 2), the SC is thick and impermeable to sedate molecules[11].

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 1125-1132 www.ijprajournal.com
ISSN: 2456-4494



Figure 2. Structure of subcutaneous layer of skin

Thus, Overcoming the SC boundary, conveying the medicine to the skin, and letting it diffuse to reach the dermal blood vessels is the most troublesome portion of TDD. The issue is that TDD is not broadly utilized in clinical hone since as it were few drugs can enter the skin. It is suggested that unused and imaginative TDD methods for upgrading skin infiltration becreated in arrange to get over this limitation. In arrange to get over the SC obstruction, a number of potential approaches, counting chemical and physical procedures, have been examined. In a 2008 distribution, Prausnitz and Langer point by point three eras of TDD frameworks. Few endeavors were made to move forward the skin penetrability of the inactive measurement shapes (transdermal patches) utilized in the to begin with era of TDD frameworks, which were aiming to provide modest, lipophilic, capable solutions. Measurement shapes that utilized iontophoresis and chemical enhancers were portion of the moment era of TDD frameworks. The methods utilized to change the SC's boundary characteristics can be isolated into two categories: chemical and physical as outlined Strategies for making strides medicate transport by means of the skin are what characterize the third era of TDD gadgets. In a 2008 ponder, Prausnitz and Langer point by point three eras of TDD systems[12].

Transdermal patches, or inactive measurement shapes, were utilized in the starting era of TDD frameworks to regulate profoundly powerful, lipophilic, minor drugs,

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 1125-1132 www.ijprajournal.com
ISSN: 2456-4494

and few endeavors were made to increment their skin porousness. Measurement shapes that utilized iontophoresis and chemical enhancers were portion of the moment era of TDD frameworks . Frameworks that depend on SC disturbance for penetration improvement instep of infiltration expansion are what characterize the third era of TDD frameworks. TDD of the third era frameworks like electroporation, microneedles (MNs), and warm removal strategies have all been appeared to be competent of conveying a distant more extensive assortment of solutions and to empower the infiltration of expansive macromolecules over the skin, such as proteins and antibodies. It is critical to note that, as Figure 3 outlines, Whereas the subcutaneous interrupt procedures to make3rd generation frameworks are generally physical strategies, the lion's share of infiltration enlargement strategies regularly connected with second-generation TDD frameworks are chemical methods. Future parts will go over more later strategies that are being inquired about for TDD frameworks that utilize a blend of chemical and physical procedures to move forward infiltration. These socalled cross breed frameworks, which are secured in more detail in the taking after areas, can be considered the "fourth era" of TDD systems[13].



Fig. 3 Methods to improve drug passes through skin

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 1125-1132 www.ijprajournal.com
ISSN: 2456-4494

IV. CHEMICAL TECHNIQUES FOR TRANSDERMAL MEDICATION ADMINISTRATION

By changing sedate and vehicle intelligent and optimizing details, chemical strategies are utilized to increment the penetrability of drugs over the SC. Due to their reasonableness, ease of generation, adaptability in plan, and capacity to empower patients to self-administer their solutions, chemical procedures are the most frequently explored technique for progressing transdermal sedate retention . Moreover, salves, gels, can be put anyplace on the skin for both immediate and long-term organization can be made by combining chemical forms with pharmaceuticals[14]. In any case, the drug's atomic weight and concentration angle generally manage the pace of medicate dissemination with this strategy, making Conveying macromolecules by means of the skin is challenging, if not inconceivable. A rate-limiting step really comes about from the necessity for a adequate concentration slope over the skin film; in arrange of medicate to discard over tissue, it have to begin with the skin's layer at tall sufficient to produce the content slope which act as a running constrain of dissemination of subcutaneous layer which slack for hour is the period of time required for sufficient of the pharmaceutical to be discharged from the transdermal dose frame (such as a transdermal fix) and construct up tissue layer for tall sufficient application to advance percutaneous infiltration. Slack time and bioavailability are the two most vital components of transdermal fix pharmacokinetics [15].Maybe the most critical impediment of the chemical procedure is the delay in pharmaceutical discharge. When an early starting of activity is required, measurement definitions with a slack period ought to not be utilized. Besides, transdermal patches' bioavailability can once in a while be lower than that of parenteral dosage shapes. For occurrence, rotigotine has a transdermal bioavailability of generally 37%. Bio accessibility Riva Stach patch fix which appeared at generally half % add up to stacking dose in another trial with 51 people with Alzheimer's infection [16].

V. CPES, OR CHEMICAL PENETRATION ENHANCERS

Chemical entrance boosts(CPEs) are broadly examine chemical strategy of changing SC's boundary characteristics . The capacity of CPEs to increment sedate atom transport through the skin has been explored [17] . They work by disturbing of arranged biological membrane , association including cavity film with biomolecule, collaboration including cytoplasmic biomolecules, disturbing, cytoplasmic triglycerides moving forward for water of subcutaneous, impacting medicate parcel factors, among other components that depend on the chemical cosmetics of CPEs . To offer assistance drugs stream through the SC, exceeding 300 chemical boosters utilized in this system. As a perfect enhancer ought to be biocompatible, nontoxic, and have consistent, unsurprising movement and length of activity. CPEs ought to encourage medicine passage into the body whereas maintaining a strategic distance from endogenous fabric misfortune (unidirectional stream) [18] . Liquor, Sulfinyl, Laurocapram, Butyro-lactum, fundamental volatiles, , grease are the most commonly utilized CPEs (Table 1) . There are a few approaches to overcome the restrictions of person chemical enhancers utilizing group of CPEs. In arrange to progress penetrability and reduce aggravation, the utilization of combinations of CPEs has been considered and authorized chemical combination appeared boost viability , security of these dose shapes in expansion to their advantageous impacts on sedate transport rates. Karande et al. appeared that systemic organization as larger particles fix utilizing a group of entrance boosters (phenyl piperazine) is doable. this found that applying two CPEs (ethanol and linoleic corrosive) on a SC demonstrate moved forward lidocaine

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 1125-1132 www.ijprajournal.com
ISSN: 2456-4494

conveyance in a synergistic way . Also, Cho et al. looked into how a entrance enhancer influenced the entrance of atenolol through evacuated rodent skin. Polyethylene glycol oleyl ether.

The vast fruitful booster when compared to glycols, greasy acids, and non-ionic surfactants. It shown a rise constantly changing associated for made strides tissue entrance[19]. CPEs have a noteworthy confinement when it include water loving large particles like affront or other biomolecules actually in spite for the fact that they frequently give 10-fold most extreme enhancements in porousness for different drugs. In reality, none of these treatments are presently accessible for buy as transdermal patches. Destitute security and productivity are the essential impediments of CPEs . The primary issues with entrance enhancers are regularly skin aggravation or maybe than a hazardous need of security [20]. Combinations of chemical patches entrance boost dynamic fixings, instruments.[21]

Table 1. Record of Chemical Penetration Enhancer with their MOA.

Chemical patch boosters	Medications	Mechanism of Action
Methylsulfynylmethyl	Cortenema,Androderm, Narcan	Disturb biomolecule, alter intertissue adaptation for subcutaneous.
1-dodecylazacycloheptan-2-one	Oruvail ,Tecfidera , Efudex	Impact on the biomolecule layer.
Pyrrolidone	Oruvail ,LignocaneHCL , Betadrenol	Modify dissolvability of subcutaneous.
Lipids	Froben, Hemangeol , Elixophyllin	Combine with lipid layer alter its covering.
Bevarages	Desmethylamitriptyline , Elixophyllin ,Lignocane HCL	Change sedate solvency in the SC Increment medicate dividing Extricate biomolecular layer.
Carbamide	Indocin ,Effexor XR, Metrogel	Disturb biomolecular layer.
Terpenoids	azidothymidine ,Tecfidera, Tofranil.	Disturb the biomolecule layer.
Solubiling Agents	Temesta ,Foscavir , Laroscorbine, Laroscorbine	Alter the compliance of the solvency.

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 1125-1132 www.ijprajournal.com
ISSN: 2456-4494

Additives	Diclofinac Potassium	Increase the influencing by decreasing the rheological characteristics. Booster loaded gels.

VI. VESICLES

The utilization of vesicular frameworks, particularly nanosized frameworks, is huge combined chemical procedures built up for impacting the subcutaneous. They can both empower controlled helpful action and progress the bioavailability of typified solutions. It was found that these frameworks might both alter systemic retention and stretch the length of medicate home in the epidermis . the squares of nearness of h_2o comes about in vesicles, which are profoundly requested congregations composed of one or more bilayers. By centering on specific zones, these frameworks can diminish poisonous quality and increment the bioavailability of medicines. Moreov, including drugs can regularly reduce issues with sedate insecurity, insolubility, and speedy corruption into vesicles.



Fig. 4.Picture of Various Types of Cellular Transport

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 1125-1132 www.ijprajournal.com
ISSN: 2456-4494

When a medicate is overseen utilizing such a conveyance framework, its clearance and tissue conveyance profile are decided by the highlights of vesicles or maybe than the physicochemical qualities of sedate particles .The medication may be included in the bilayer or encased in the insides depth, depending on its physicochemical properties . Subsequently, hydrophilic, lipophilic, and amphiphilic drugs can be stacked into vesicles for transdermal conveyance . conveyance and embodiment drugs[22]. Lipid Vesicles found in 1965, biomolecular materials utilized broadly drug, Because phospholipids and greasy acids are normally plenteous in cell layers , Lipid layers structures similar to watery inner surroundings are need for biowastes. When conveyed in watery conditions, they arrange themselves so that the greasy corrosive bunches confront one another and the polar head bunches confront outward into the fluid locale, producing round, vesicle-like structures that take after lamellar sheets . Phospholipids, surfactants, and an inward watery compartment encased in a lipid bilayer that can typify hydrophilic and lipophilic drugs make up deformable liposomes As edge activators, surfactants increment vesicle deformability and destabilize lipid bilayers . Deformable liposomes are alluded to as adaptable, versatile, ultradeformable, and ultra-flexible liposomes[23].

Deformable liposomes have been demonstrated to have superior helpful impacts than ordinary liposomes since they can saturate more profound into epidermal layers[24].Transfersomes make up the starting era of flexible liposomes. Exchange somes, some of the time alluded the higher versatile niosomes with boundary booster and a phospholipid spine . The essential cause of the vesicles' flexibility is the boundary booster, debilitates their biolayer and makes strides their adaptability and deformability. The thin-film hydration approach is regularly utilized to plan transfersomes. Transfersomes' progressed skin infiltration is for the most part due to their deformability and the osmotic slope over the application location [25].

Elastic Vesicles , flexible liposomes utilized by its framework, concurring to Duangjit et al. since they illustrated essentially more noteworthy meloxicam skin infiltration than liposomes. Analysts regularly utilize transfersomes as transdermal conveyance vehicles as antineoplastic medicines, NSAIDS, anesthesia, affront, and since of their ultra-deformability. In spite of their numerous benefits, transfersomal conveyance frameworks are costly, chemically unsteady, and phospholipid quality is another critical thought . In later decades, researchers working in the field of sedate conveyance frameworks have gotten to be interested in the advancement of surfactant vesicles, or niosomes, as a way to move forward TDD. corticosteroids . similar to niosomes which can flow water loving and water hating medicaments whereas moreover conveying for aim [26].

VII. CONCLUSION

Due to its preferences, counting noninvasiveness, self-administration, and the capacity to give drugs reliably at predefined and directed rates, TDD is presently getting to be more and more prevalent as a conveyance strategy for a number of disarranges. Subsequently, TDD innovation is picking up footing in the pharmaceutical division. Drugs have been conveyed to the skin by means of a assortment of chemical transdermal conveyance stages, counting nanoparticles, liposomes, nanocrystals, niosomes, and nanoemulsions. Be that as it may, A stratum Cornea is a anticipates water loving particles or large particles entering intaglio tissue, is occasionally experienced by these conveyance stages. It has been illustrated that physical increase methods make strides sedate conveyance of its site of action, empowering of organization for assortment of doses, which that they can frequently challenging for regulate utilizing strategies for

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 1125-1132 www.ijprajournal.com
ISSN: 2456-4494

making strides chemical infiltration, such macromolecules. The binding of applicants which are utilizing blend of beneath the skin gadgets or different dosage contained details is one of the challenges that still require to be settled. Also, certain rising transdermal approaches, such microneedles, require administrative supervision over fabricating and scaled-up GMP.

REFERENCES

[1]. Sahoo D, Bandaru R, Samal SK, Naik R, Kumar P, Kesharwani P, Dandela R. Oral drug delivery of nanomedicine. In: Theory and Applications of Nonparenteral Nanomedicines. Academic press 2021,vol. 27 (3), 181-207.

[2]. He M, Zhu L, Yang N, Li H, Yang Q. Recent advances of oral film as platform for drug delivery. Kaur G, Arora M, Ravi Kumar MNV; Book: Oral Drug Delivery Technologies a decade development . J Pharmacol. Exp. Ther. 2021 ; vol.370 (2) ; 529-543.

[3]. Raza A, Hayat U, Bilal M, Iqbal HMN, Wang J-Y. Zein-based micro- and nano-constructs and biologically therapeutic cues with multi-functionalities for oral drug delivery systems. J drug delivery science technology, 2020 ; vol.58(5) ; 101-818 .

[4]. Gulati N, Gupta H. Parenteral drug delivery: A review. Recent Pat Drug Deliv Formul, 2011; vol.226(5): 133–145.

[5]. Nagarsenkar MS, Dhawan VV. Parenteral preparations. In: Remington, 23rd ed. Academic Press, 2021; vol.5(1) ; 577–603.

[6]. Pastore MN, Kalia YN, Horstmann M, Roberts MS. Transdermal patches: History, development and pharmacology. Br J Pharmacol, 2015; vol.172(6): 2179–2209.

[7]. Prausnitz MR, Langer R. Transdermal drug delivery. Nat Biotechnol, 2008; vol.26(9): 1261–1268.

[8]. Sharma G, Alle M, Chakraborty C, Kim JC. Strategies for transdermal drug delivery against bone disorders: A preclinical and clinical update. J Control Release, 2021; vol.336(9): 375–395.

[9]. Jeong WY, Kwon M, Choi HE, Kim KS. Recent advances in transdermal drug delivery systems: A review. Biomater Res, 2021; vol.25(24)224-232.

[10]. Jung JH, Jin SG. Microneedle for transdermal drug delivery: Current trends and fabrication. J Pharm Investig, 2021; vol.51(36): 503–517.

[11]. Khoury R, Rajamanickam J, Grossberg GT. An update on the safety of current therapies for Alzheimer's disease: Focus on rivastigmine. Ther Adv Drug Saf, 2018; vol.9(15): 171– 178.

[12]. Tsuno N, Mori T, Ishikawa I, Bando N, Park H, Matsumoto Y, et al. Efficacy of rivastigmine transdermal therapy on low food intake in patients with Alzheimer's disease.

[13]. GeriatrGerontol Int, 2019; vol.19(6): 571–576.

[14]. Alkilani AZ, McCrudden MT, Donnelly RF. Transdermal drug delivery: Innovative pharmaceutical developments based on disruption of the barrier properties of the stratum corneum. Pharmaceutics, 2015; vol.7(24): 438–470.

[15]. Bajaj S, Whiteman A, Brandner B. Transdermal drug delivery in pain management. Contin EducAnaesth Crit Care Pain, 2011; vol.11(7): 39–43.

[16]. Abrams LS, Skee DM, Natarajan J, Wong FA, Anderson GD. Pharmacokinetics of a contraceptive patch containing norelgestromin and

ethinylestradiol at four application sites. Br J Clin Pharmacol, 2002; vol.53(18): 141–146.

[17]. Frampton JE. Rotigotine transdermal patch: A review in Parkinson's disease. CNS Drugs, 2019; vol.33(24): 707–718.

[18]. Schaer DH, Buff LA, Katz RJ. Sustained antianginal efficacy of transdermal nitroglycerin patches using an overnight nitrate-free interval. Am J Cardiol, 1988; 61(13): 146–150.

[19]. Vartan CM. Buprenorphine transdermal patch: An overview for use in chronic pain. US Pharm, 2014; vol.10(16)230-245.

[20]. Tanida N, Akiyama K, Terahara T. Pharmacological profile and clinical efficacy of transdermal patch containing emedastinedifumarate. Nihon Yakurigaku Zasshi, 2018; vol.152(25): 246–255.

[21]. Birks JS, Grimley Evans J. Rivastigmine for Alzheimer's disease. Cochrane Database Syst Rev, 2015; vol.4(27): 1000-1191.

[22]. Westhoff CL, Reinecke I, Bangerter K, Merz M. Impact of body mass index on suppression of follicular development and ovulation using a transdermal patch. Contraception, 2014; vol.90(50): 272–279.

[23]. Yousef H, Alhajj M, Sharma S. Anatomy, Skin (Integument), Epidermis. StatPearls, 2022; vol.25(2) 157-165.

[24]. Kahraman E, Kaykın M, Şahin Bektay H, Güngör S. Recent advances on topical application of ceramides to restore barrier function of skin. Cosmetics, 2019; 6(52) 125314.

[25]. Bouwstra JA, Groenink HWW, Kempenaar JA, Romeijn SG, Ponec M. Water distribution and natural moisturizer factor content in human skin equivalents regulated by relative humidity. J Investig Dermatol, 2008; vol.128(15): 378–384.

[26]. Lai-Cheong JE, McGrath JA. Structure and function of skin, hair and nails. Medicine, 2021; vol49(32): 337–342.

[27]. Al-Akayleh F, Adwan S, Khanfar M, Idkaidek N, Al-Remawi M. A novel eutectic-based transdermal delivery system for risperidone. AAPS PharmSciTech, 2020; vol.22(4)345370.