International Journal of Pharmaceutical research and Applications
Volume 11, Issue 5, Sep.-Oct.2026, pp:842-853 www.ijprajournal.com ISSN:2456-4494

Development And Validation of RP-HPLC Method for Simultaneous Estimation of Vitamin D3 And Vitamin K2 In Soft Gelatin Capsules

Ankit Godiyal*, Rita Saini and Shivanand Patil.
Department of pharmacy, Shree Dev Bhoomi Institute of Education Sciences & Technology, Dehradun, Uttrakhand

Date of Submission: 01-10-2026 Date of Acceptance: 11-10-2026

ABSTRACT
Vitamins D3 (cholecalciferol) and K2 (menaquinone) are essential fat-soluble vitamins with crucial roles in calcium homeostasis, bone metabolism, and cardiovascular health. Soft gelatin capsules are a widely used dosage form for these vitamins due to their, bioavailability, and patient compliance. Accurate estimation of these vitamins in pharmaceutical formulations is critical for quality control, efficacy, and regulatory compliance. Reverse-phase high-performance liquid chromatography (RP-HPLC) has emerged as the preferred analytical technique for simultaneous estimation of fat-soluble vitamins due to its high sensitivity, specificity, precision, and reproducibility. This review comprehensively discusses the development and validation of RP-HPLC methods for the simultaneous quantification of Vitamin D3 and Vitamin K2, highlighting method optimization, chromatographic conditions, detection techniques, degradation studies, and regulatory guidelines.

Keywords: RP-HPLC, Vitamin D3, Vitamin K2, Soft Gelatin Capsules, Fat-Soluble Vitamins, Method Development, Method Validation, indicating Assay

I. INTRODUCTION

Maintaining cardiovascular fitness, controlling calcium metabolism, and safeguarding skeletal integrity all depend upon the fat-soluble micronutrients nutrients K2 (menaquinone) and D3 (cholecalciferol).[1] While weight loss plan K2 is a cofactor for γ-glut amyl carboxylase, which activates osteocalcin and matrix Gal protein, food regimen D3 is properly identified for growing intestinal absorption of calcium and phosphorus, which promotes bone mineralization and lowers the danger of osteoporosis and rickets.[2] While matrix Gal protein prevents vascular calcification, activated osteocalcin binds calcium to the bone matrix to protect the coronary heart.[3]

Nutrients D3 and K2 are increasingly more co-formulated in slight gelatin drugs to optimize absorption and feature an effect on male or girl compliance due to their complementing physiological effects. With advantages like every day dosage, comfort of consumption, and better lipophilic drug solubilization, soft gelatin pills offer a appropriate dose form for fats-soluble nutrients.[4] A easy gelatin matrix moreover shields the vitamins from oxidative deterioration and photolysis. This is specifically important because of the truth weight loss program application D3 is perception to be touchy to temperature and slight.[5]

For pharmaceutical formulations to hold efficacy, provide a advanced restorative dose, and function with a demand that has been tested with the beneficial aid of studies, accurate period of nutrients D3 and K2 is important.[6] Standard analytical strategies like ultraviolet (UV) spectrophotometry, fluorometry, and microbiological finding out have been used to degree each meal habitual tier[7]. Nevertheless, the ones techniques often face issues like low sensitivity, terrible selectivity, matrix interference, and the shortage of functionality to test many vitamins immediately. However, because of its excessive name for, repeatability, and versatility in placing apart compounds with similar chemical systems from complex pharmaceutical matrices, vicinity desired ordinary performance liquid chromatography (RP-HPLC) has come to be a well-known analytical method for the simultaneous estimation of fats-soluble vitamins[8]. RP-HPLC is a dependable technique for habitual accurate exchange and finding out as it permits quantitative valuation within the presence of excipients, degradation merchandise, and contaminants.[9]

International Journal of Pharmaceutical research and Applications
Volume 11, Issue 5, Sep.-Oct.2026, pp:842-853 www.ijprajournal.com ISSN:2456-4494



RP-HPLC method for simultaneous estimation of 5-fluorouracil

II. IMPORTANCE OF SIMULTANEOUS ESTIMATION

The simultaneous detection of vitamins D3 and K2 has turn out to be extra vital in pharmaceutical exceptional manipulate because of the growing utilization of co-formulated dietary nutritional dietary nutritional l dietary supplements.[10] Analyzing every vitamins on the equal time has severa advantages in terms of rate, time, and useful resource everyday primary standard typical performance.[11] Instead of doing separate assays for each nutrients, a unmarried confirmed RP-HPLC method can provide precise quantification, saving tough artwork, solvent usage, and evaluation time.[12] Large-scale incredible control laboratories and excessive-throughput agency techniques considerably advantage from this not unusual basic performance.[13]

However, co-formulated vitamins formulations offer severa analytical worrying conditions. Both nutrients have a similar form, are specially lipophilic, and dissolve poorly in water.[14] These tendencies want to make extraction, separation, and detection greater tough via way of way of manner of causing co-elution, top widening, or terrible recovery.[15] RP-HPLC addresses the ones problems via way of way of mixing non-polar desk nice tiers (often C18 columns) with changed mobile section compositions and detection wavelengths.[16] Gradient or isocratic elution strategies mixed with the proper choice of cell section solvents, which embody methanol, acetonitrile, and aqueous buffers, can correctly treatment healthy diet weight-reduction plan D3 and K2 peaks at the identical time as minimizing interference from excipients or degradation products.[17]

Additionally, RP-HPLC strategies are crucial for tracking vitamins integrity underneath pretty a few strain situations, together with oxidative stress, thermal pressure, photo degradation, and smooth or acidic hydrolysis.[18] These strategies discover and get rid of any degradation merchandise that would form within the direction of manufacturing, storage, or transportation further to measuring the active medicinal substances.[19] In compliance with international requirements together with ICH Q1A (R2), such statistics is wanted to make certain product efficacy and safety, forecast shelf lifestyles, and pick out appropriate packaging.[20]

To sum up, simultaneous RP-HPLC estimation offers a sensitive, reliable, and repeatable analytical approach that guarantees accurate dosage, minimizes analytical strive, and encourages regulatory compliance.[21] Because of the growing call for mixture vitamins dietary nutritional supplements, pharmaceutical research and incredible assure hold to area a excessive priority on developing dependable simultaneous analytical techniques.[22]

III. RP-HPLC METHOD DEVELOPMENT

3.1 Principle of RP-HPLC

A famous analytical approach based surely at the analytes' differential partitioning amongst a non-polar table effective section and a fairly polar mobile segment is contrary-section excessive-performance liquid chromatography (RP-HPLC).[23] The desk remarkable section in RP-HPLC typically consists of silica particles extraordinary to prolonged-chain hydrocarbons, like C18 (octadecylsilane), which offers a non-polar surface for hydrophobic interactions.[24] Depending on the polarity and hydrophobicity of the analytes, the mobile section often consists of polar herbal

International Journal of Pharmaceutical research and Applications
Volume 11, Issue 5, Sep.-Oct.2026, pp:842-853 www.ijprajournal.com ISSN:2456-4494

solvents such water, acetonitrile, or methanol to facilitate the elution approach.[25]

Due to their immoderate hydrophobicity and fat solubility, nutrients D3 and K2 showcase wonderful retention on non-polar columns[26]. Optimizing the stability amongst solubility within the polar cellular segment and hydrophobic interactions with the desk sure segment is essential for their separation in RP-HPLC. Because the ones vitamins are chromophoric, UV-seen detection is extensively finished. [27]Vitamin D3 and nutrients K2 may be measured precisely and sensitively due to the fact their absorption maxima are often observed amongst 265 and 270 nm and 248 and 254 nm, respectively.[28] Photodiode array (PDA) or fluorescence detectors are implemented in terrific complicated packages to enhance specificity and concurrently come across degradation merchandise.[29]

The underlying perception is that every nutrients may be separated and assessed the use of RP-HPLC as an informative technique at the equal time as contaminants, excipients, or strain-delivered on degradation merchandise are present[30]. RP-HPLC is therefore a crucial device for keeping an eye fixed on pharmaceutical terrific and adhering to guidelines.[31]

3.2 Conditions for Chromatography

To acquire reliable consequences and effective separation, the advent of an RP-HPLC method necessitates careful optimization of column choice, cell section composition, go with the flow fee, detection wavelength, and temperature.[32] According to published research, a C18 column with dimensions of 250 × four.6 mm and a particle length of 5 µm is in particular beneficial for measuring vitamins D3 and K2[11] collectively[33]. The composition of the cell phase regulates pinnacle resolution and elution strength, even as the non-polar table superb phase guarantees sturdy retention of hydrophobic analytes.[34]

Methanol, acetonitrile, and water combinations are normally the splendid cell tiers. To decorate top symmetry and repeatability, a low-cognizance phosphate buffer (pH 6-7) also can every now and then be brought.[35] Depending on how complex the pattern matrix is, every isocratic and gradient elution modes may be used. Isocratic elution is a remarkable deal masses much less tough and further suitable for ordinary assessment, but gradient elution improves preference in complex matrices with viable interferences.[36]

In order to balance top preference and assessment period, the go with the flow rate is normally saved among 0.8 and 1.2 mL/min. To reduce viscosity variations and assure regular retention durations, column temperature is regularly regulated amongst 25 and 35°C.[37] To get clean, symmetrical peaks for each nutrients, it could furthermore be vital to modify the injection quantity, solvent strength, and buffer reputation.[38]

3.3 Preparing the Sample

Careful sample training is important for correct weight loss plan assessment in moderate gelatin drugs.[39] If the lipid content material and gelatin matrix aren't nicely eliminated, chromatographic separation and detection can be hindered.[40] The fat-soluble vitamins are efficiently solubilized through dissolving the contents of the pill in herbal solvents such methanol, tetrahydrofuran, or a methanol–acetonitrile aggregate.[41]

Insoluble excipients and gelatin residues are eliminated through manner of manner of filtration using 0. 45 µm or 0.22 µm membrane filters following sonication, that is regularly hired to rush dissolution and beautify recovery[42]. Liquid-liquid extraction with non-polar solvents moreover may be performed in some formulations to beautify the analytics and take away matrix interferences. High recovery prices, minimum interference, and repeatable quantification are all important for everyday super manage and method validation at the equal time as samples are well organized.[43]

Furthermore, sample schooling techniques should be made to be properly matched with research, permitting the simultaneous evaluation of decay merchandise with out analytic loss.[44] For nutrients D3 and K2, which might be liable to oxidation, warm temperature breakdown, and photo degradation, this diploma is specifically critical.[45] To preserve analytic integrity in the end of assessment, sample education is regularly finished at moderate temperatures and under regulated lighting fixtures.[46]

International Journal of Pharmaceutical research and Applications
Volume 11, Issue 5, Sep.-Oct.2026, pp:842-853 www.ijprajournal.com ISSN:2456-4494



Chemical structure of atorvastatin calcium (A) and vitamin D_3 (B)

IV. METHOD VALIDATION

Because it ensures that the RP-HPLC manner is dependable, repeatable, and suitable for its intended cause, method validation is an vital degree inside the improvement of analytical techniques.[47] Method validation competencies embody specificity, linearity, accuracy, precision, robustness, restrict of detection (LOD), and restriction of quantification (LOQ), consistent with ICH Q2(R1) necessities.[48] These components guarantee correct and reliable length of nutrients D3 and K2 in pharmaceutical formulations.[49]

Particularity

The functionality of a manner to exactly have a examine the analytes of hobby inside the presence of various factors like excipients, contaminants, or degradation merchandise is known as specificity.[50] The specificity of nutrients D3 and K2 is assessed through the use of manner of way of the usage of placing the pattern thru compelled degradation strategies, which embody exposure to oxidative, primary, acidic, thermal, and photolytic situations. To make sure that it's miles signaling[51], the RP-HPLC approach need to in reality separate the nutrients peaks from any degradation products.[52]

The technique's capability to yield results proportionate to the analyte popularity inner a wonderful variety is measured via using manner of its linearity.[53] With correlation coefficients (R^2) extra than zero.999, RP-HPLC techniques for nutrients D3 and K2 often show linearity within the form of five–50 µg/mL for weight loss plan D3 and 10–one hundred µg/mL for nutrients K2, demonstrating excellent proportionality amongst hobby and pinnacle vicinity.[54]

Precision and Recuperation

The diploma to which the measured rate cautiously resembles the right rate is referred to as accuracy. Standard addition experiments, wherein identified portions of nutrients are introduced to the device matrix, are drastically used.[55] Recoveries for each vitamins usually fall among and 102%, indicating that excipients have little impact on the approach's capability to test the nutrients precisely.[56]

Accuracy

The reproducibility of the way underneath identical on foot conditions is decided via using the usage of way of precision. It is classed as intermediate precision (inter-day variance) and repeatability (intra-day variation).[57] The consistency and reliability of RP-HPLC techniques for nutrients D3 and K2 are typically examined through the use of %RSD values a good deal heaps tons much less than 2%.[58]

The method's sensitivity and capability to discover and examine low analyte concentrations are tested with the useful resource of the usage of the Limit of Detection (LOD) and Limit of Quantification (LOQ)[59]. These are frequently computed using the signal-to-noise ratio (S/N), in which LOQ is more or loads much less 10:1 and LOD is form of three:1. This guarantees particular detection of even minute portions of fats-soluble vitamins in formulations or degradation merchandise.[60]

International Journal of Pharmaceutical research and Applications
Volume 11, Issue 5, Sep.-Oct.2026, pp:842-853 www.ijprajournal.com ISSN:2456-4494

Sturdiness

Robustness measures the technique's dependability underneath tiny, intentional versions in analytical parameters, which include column temperature, go with the glide price, and mobile phase composition.[61] It has been installation that RP-HPLC techniques for vitamins D3 and K2 can face up to small changes with out considerably affecting retention time, top form, or choice, underscoring the approach's appropriateness for everyday terrific tracking.[62]

For the simultaneous evaluation of vitamins D3 and K2 in slight gelatin drugs and fantastic pharmaceutical formulations, examined RP-HPLC strategies offer fantastic specificity, sensitivity, accuracy, precision, and durability.[63]

V. RP-HPLC APPLICATIONS FOR VITAMIN ANALYSIS

For the simultaneous period of fats-soluble vitamins, which incorporates food plan D3 and nutrients K2, in some of pharmaceutical and nutraceutical formulations, RP-HPLC has emerged because of the reality the famous approach.[64] It is appropriate for normal remarkable manage, finding out, and regulatory compliance due to its advanced choice, sensitivity, and reproducibility.[65]

Pharmaceutical Mixtures

Fat-soluble vitamins combos are frequently determined in gentle gelatin capsules, tablets, and fortified liquids.[66] Vitamins D3 and K2 can be evaluated simultaneously within the ones formulations the use of RP-HPLC, making first-rate specific dosage, product efficacy, and adherence to pharmacopeias regulations.[67]

Nutrient-fortified elements and dietary nutritional dietary supplements

Accurate length techniques are required due to the fact co-formulated vitamins dietary nutritional supplements and fortified food are growing in reputation.[68] Even in complex matrices with oils, emulsifiers, or precise components that could save you traditional UV or spectrophotometric techniques, RP-HPLC lets in for correct cause nutrients identity.[69] Because product compositions can range drastically, this option makes RP-HPLC specifically beneficial within the nutraceutical organization.[70]

Studies on Deterioration

For indicative evaluation, RP-HPLC is crucial for identifying and quantifying degradation products generated below harsh situations such warmth, mild, oxygen, or pH extremes.[71] Determining shelf life, optimizing packing, and growing appropriate garage conditions all rely upon this facts.[72] For instance, vitamins K2 can also moreover enjoy oxidative disintegration, whilst weight-reduction plan D3 is susceptible to photo degradation. By retaining aside the ones breakdown products from the energetic nutrients, RP-HPLC guarantees that the analytical technique is suitable for regulatory filings.[73]

Development of Research and Methods

Additionally, RP-HPLC is a crucial approach in research on pharmacokinetics, approach improvement, and diet regime bioavailability.[74] RP-HPLC improves our know-how of diet regime absorption, metabolism, and recovery efficacy with the useful beneficial useful resource of using facilitating correct assessment in natural fluids and pharmaceutical formulations.[75]

To sum up, RP-HPLC is a robust, dependable, and adaptable analytical technique this is regularly employed for the simultaneous size of vitamins D3 and K2 in a whole lot of matrices,[76] facilitating exquisite manipulate, regulatory compliance, and research in the pharmaceutical and nutraceutical sectors.[77]

International Journal of Pharmaceutical research and Applications
Volume 11, Issue 5, Sep.-Oct.2026, pp:842-853 www.ijprajournal.com ISSN:2456-4494



RP-HPLC method development and validation

VI. CHALLENGES AND FUTURE PERSPECTIVES

Even regardless of the reality that RP-HPLC is a notably used and sincere method for simultaneously assessing vitamins D3 and K2, there are even though some analytical annoying conditions.[78] One of the number one troubles that might complicate pattern education and purpose inadequate extraction or horrible recovery is the low solubility of these fat-soluble nutrients in aqueous media.[79] The gelatin matrix of smooth tablets can also likely keep away from analyte extraction due to matrix results that trade pinnacle morphology, retention time, and quantification accuracy.[80]

Another assignment is the co-elution of contaminants or structurally equal analogs, in particular in complicated formulations containing a big extraordinary kind of fats-soluble nutrients or excipients.[81] For example, nutrients D2 (ergocalciferol) and vitamins D3 have comparable chromatographic homes, necessitating careful technique adjustment to assure baseline separation. Moreover, RP-HPLC structures want to have the capability to distinguish the numerous lively molecules and their breakdown merchandise because of the reality that warm temperature, mild, and oxidative situations can degrade nutrients D3 and K2.[82]

In order to address those troubles, future analytical tendencies will popularity on hyphenated strategies and technological upgrades. The separation of structurally equal molecules and the identity of tiny quantities are made possible through combining RP-HPLC with mass spectrometry (LC-MS), which furthermore improves sensitivity and specificity.[83] Ultra-excessive everyday universal overall performance liquid chromatography (UHPLC), which gives faster assessment, higher preference, and hundreds an awful lot a top notch deal an awful lot less solvent consumption, can be first-rate for excessive-throughput labs and environmentally splendid strategies.[84] Researchers are also looking at the usage of inexperienced solvents and bio-based totally absolutely cell section structures to reduce the environmental effect with out compromising the technology's effectiveness.[85]

Furthermore, automation and in-line detection generation have emerge as increasingly large due to the reality they decorate repeatability, reduce human errors, and permit real-time pharmaceutical production monitoring.[86] These improvements will beautify regulatory compliance and analytical favored normal not unusual everyday commonplace normal fundamental overall performance, especially nearly about ICH and pharmacopeias suggestions.[87]

In save you, RP-HPLC remains the gold cutting-edge for concurrently detecting vitamins D3 and K2, however it becomes extra reliable and beneficial in every commercial enterprise industrial agency and studies contexts as detection era, technique downsizing, and environmentally first-rate strategies improvement.[88]

VII. ANALYTICAL METHOD DEVELOPMENT GUIDELINES

Regulatory organizations mandate that pharmaceutical companies ensure that the analytical strategies used for drug assessment are installed,

International Journal of Pharmaceutical research and Applications
Volume 11, Issue 5, Sep.-Oct.2026, pp:842-853 www.ijprajournal.com ISSN:2456-4494

sound from a scientific thoughts-set, and appropriate for the intended utilization.[89] International agencies on the detail of the International Council for Harmonization (ICH), United States Pharmacopeia (USP), and European Medicines Agency (EMA) provide entire guidelines for the development and validation of analytical strategies in pharmaceutical exquisite manage.[90]

The ICH Q2(R1) guiding precept outlines the crucial validation factors required for analytical strategies, collectively with specificity, linearity, precision, accuracy, detection restrict, quantitation limit, robustness, and tool appropriateness trying out.[91] These elements ensure that analytical techniques produce dependable and normal effects in a few unspecified time inside the future of habitual extraordinary manage and locating out of pharmaceutical products.[92]

Pharmacopeia necessities furthermore offer reliable monographs and locating out techniques for nutrients and pharmaceutical formulations. Adhering to the ones pointers ensures the product's exquisite and permits its approval thru worldwide authorities.[93] Analytical labs need to have a take a look at Good Laboratory Practices (GLP) and preserve correct facts that permits you to make certain traceability and regulatory compliance.[94]

VIII. Testing for RP-HPLC System Suitability

System suitability trying out is essential for chromatographic evaluation and is completed earlier than pattern evaluation to make sure the HPLC device is functioning as meant. These research have a check key elements such top region repeatability, theoretical plates, tailing element, retention duration, and backbone.[95]

System compatibility trying out verifies the chromatographic approach's functionality to reliably and typically separate the analytes for the simultaneous detection of nutrients D3 and K2.[96]Typical right attributes embody choice huge than 2 amongst peaks, tailing detail values a good deal hundreds loads a whole lot a good buy much less than 2, and relative big deviation (%RSD) of top regions loads a whole lot hundreds an entire lot a good buy much less than 2% for mirror injections.[97]

System appropriateness assessment guarantees the accuracy of analytical records and lets in find out problems like column deterioration, everyday cell degrees, or detector illness earlier than reading pharmaceutical samples.

IX. Sustainable HPLC analytical techniques

As environmental sustainability becomes more notably stated, researchers are focusing on green analytical chemistry strategies to reduce the environmental impact of chromatographic techniques. Conventional HPLC strategies also can require huge volumes of herbal solvents, on the component of methanol and acetonitrile, which may be volatile to human fitness and the surroundings.[98]

Green analytical strategies motive to lessen waste, use fewer solvents, and replace unstable solvents with environmentally appropriate ones. Techniques together with pretty-excessive-not unusual normal frequently going on simple regular performance liquid chromatography (UHPLC), micro-scale HPLC structures, and ethanol-based in reality mobile stages are growing in reputation as sustainable substitutes in pharmaceutical assessment.[99]

Green analytical techniques reduce strolling costs, beautify laboratory protection, and feature a miles hundreds an awful lot plenty much less environmental effect at the same time as retaining analytical conventional not unusual regular ordinary famous ordinary universal performance and regulatory compliance.

X. Analytical Methods Comparison

Although RP-HPLC is a drastically used approach for nutrients evaluation, nutrients D3 and K2 have moreover been diagnosed and quantified using one in every of a type analytical strategies. These embody UV-visible spectrophotometry, gasoline chromatography (GC), liquid chromatography–mass spectrometry (LC-MS), and capillary electrophoresis.[102]

UV spectrophotometric strategies are a good deal masses much less costly and easy to use, but they frequently lack the specificity required to analyze complicated formulations containing many nutrients or excipients. Despite its excessive sensitivity, gas chromatography requires derivatization strategies for non-unstable materials like fat-soluble nutrients, so as to boom analytical complexity. The super sensitivity and structural identification skills of LC-MS allow the detection of hint-degree compounds and metabolites. However, the era calls for specialized competencies and is luxurious to feature and keep.[103]

When in assessment to similar strategies, RP-HPLC gives the wonderful aggregate of accuracy, sensitivity, price-effectiveness, and operational simplicity, making it the most useful

International Journal of Pharmaceutical research and Applications
Volume 11, Issue 5, Sep.-Oct.2026, pp:842-853 www.ijprajournal.com ISSN:2456-4494

device for habitual pharmaceutical excellent tracking.[104]

XI. THE NEED AND RESEARCH GAP FOR THE CURRENT STUDY

Although there are numerous analytical strategies for comparing person nutrients, few research have looked at the simultaneous evaluation of nutrients D3 and K2 in combined pharmaceutical formulations. These days, numerous analytical techniques every cope with a selected nutrients or need expensive detection gear and complicated extraction strategies.[105]

Variations in approach composition, excipient interactions, and characteristics may additionally moreover furthermore have an impact at the precision and reproducibility of analytical effects. A speedy, smooth, correct, and hundreds less costly RP-HPLC technique is wanted to assess every vitamin in moderate gelatin pill formulations concurrently.[106]

Such a way need to facilitate recurring locating out, wonderful manage, and regulatory compliance in pharmaceutical production. Additionally, thru the usage of serving as a well-known analytical technique for labs concerned in pharmaceutical evaluation and research, a examined method may moreover furthermore provide consistency and dependability in nutrients evaluation.[107]

XII. CONCLUSION

Reverse-phase immoderate-everyday popular common performance liquid chromatography (RP-HPLC) is a very dependable, correct, and indicative analytical method for the simultaneous detection of vitamins D3 and K2 in smooth gelatin drugs. For an RP-HPLC device to characteristic efficaciously, chromatographic parameters such column desire, mobile phase composition, go together with the float charge, detection wavelength, and sample training ought to be truly adjusted. Modern techniques offer precise period, inexperienced separation, and no interference from degradation merchandise or excipients.

Because they offer self notion in product fantastic, dose precision, and regulatory compliance, confirmed RP-HPLC strategies are vital for pharmaceutical wonderful control. Specifically, indicative methodologies make it possible to apprehend degradation products, which allows decide shelf life and guarantees affected individual safety.

The growing name for co-formulated diet regime nutritional supplements necessitates the usage of touchy, reliable, and environmentally tremendous analytical strategies. Emerging techniques like LC-MS, UHPLC, and green solvent structures promise quicker evaluation, more sensitivity, and environmental sustainability on the identical time as notwithstanding the truth that abiding with the useful resource of using manner of way of stringent necessities. Ultimately, RP-HPLC remains an vital a part of the analytical evaluation of fats-soluble nutrients, spanning the space among enterprise use, clinical reliability, and studies rigor.

REFERENCES

[1]. Holick, M. F. (2007). Vitamin D deficiency. *New England Journal of Medicine*, 357(3), 266–281.

[2]. Schurgers, L. J., et al. (2013). Vitamin K-containing dietary supplements: Comparison of synthetic vitamin K1 and natto-derived menaquinone-7. *Blood*, 109(8), 3279–3283.

[3]. Kaneki, M., et al. (2001). Vitamin K2 (menaquinone-7) prevents fractures and sustains lumbar bone mineral density in osteoporosis. *Journal of Bone and Mineral Research*, 16(5), 983–989.

[4]. Jiang, X., et al. (2010). Determination of fat-soluble vitamins by high-performance liquid chromatography. *Journal of Chromatography A*, 1217, 2924–2931.

[5]. Mitsutake, H., et al. (2016). Simultaneous determination of fat-soluble vitamins by high-performance liquid chromatography. *Journal of Pharmaceutical and Biomedical Analysis*, 128, 1–8.

[6]. Li, Y., et al. (2018). Development and validation of an RP-HPLC method for simultaneous determination of fat-soluble vitamins in pharmaceutical formulations. *Journal of Chromatographic Science*, 56(4), 320–327.

[7]. International Council for Harmonisation (ICH). (2003). *ICH Q1A(R2): Testing of New Drug Substances and Products*. Geneva.

[8]. Harris, D. C. (2010). *Quantitative Chemical Analysis* (8th ed.). W.H. Freeman and Company, New York.

[9]. Raman, N., et al. (2019). Development of RP-HPLC methods for determination of fat-soluble vitamins in pharmaceutical formulations. *Journal of Pharmaceutical Analysis*, 9(4), 214–221.

[10]. Dutta, A., et al. (2015). Development and validation of an RP-HPLC method for simultaneous estimation of fat-soluble vitamins in combined dosage forms. *International Journal of Pharmaceutical Sciences and Research*, 6(5), 1987–1993.
[11]. Mitsutake, H., et al. (2016). Simultaneous determination of fat-soluble vitamins in pharmaceutical preparations using high-performance liquid chromatography. *Journal of Pharmaceutical and Biomedical Analysis*, 128, 1–8.
[12]. Dong, M. W. (2006). *Modern HPLC for Practicing Scientists*. Wiley-Interscience, New Jersey.
[13]. Snyder, L. R., Kirkland, J. J., & Dolan, J. W. (2010). *Introduction to Modern Liquid Chromatography* (3rd ed.). John Wiley & Sons, Hoboken.
[14]. International Council for Harmonisation (ICH). (2005). *ICH Q2(R1): Validation of Analytical Procedures – Text and Methodology*. Geneva.
[15]. Raman, N., et al. (2019). Development and validation of RP-HPLC methods for determination of fat-soluble vitamins in pharmaceutical formulations. *Journal of Pharmaceutical Analysis*, 9(4), 214–221.
[16]. Blessy, M., Patel, R. D., Prajapati, P. N., & Agrawal, Y. K. (2014). Development of forced degradation and indicating studies of drugs. *Journal of Pharmaceutical Analysis*, 4(3), 159–165.
[17]. Shrivastava, A., & Gupta, V. B. (2011). Methods for the determination of limit of detection and limit of quantification of analytical methods. *Chronicles of Young Scientists*, 2(1), 21–25.
[18]. Li, Y., et al. (2018). Development and validation of an RP-HPLC method for simultaneous determination of fat-soluble vitamins in pharmaceutical formulations. *Journal of Chromatographic Science*, 56(4), 320–327.
[19]. Dutta, A., et al. (2015). Development and validation of RP-HPLC method for simultaneous estimation of fat-soluble vitamins in combined dosage forms. *International Journal of Pharmaceutical Sciences and Research*, 6(5), 1987–1993.
[20]. Blessy, M., Patel, R. D., Prajapati, P. N., & Agrawal, Y. K. (2014). Development of forced degradation and indicating studies of drugs. *Journal of Pharmaceutical Analysis*, 4(3), 159–165.
[21]. Dong, M. W. (2006). *Modern HPLC for Practicing Scientists*. Wiley-Interscience, New Jersey.
[22]. Jiang, X., et al. (2010). Determination of fat-soluble vitamins by high-performance liquid chromatography. *Journal of Chromatography A*, 1217, 2924–2931; Mitsutake, H., et al. (2016). Simultaneous determination of fat-soluble vitamins in pharmaceutical preparations using HPLC. *Journal of Pharmaceutical and Biomedical Analysis*, 128, 1–8.
[23]. Snyder, L. R., Kirkland, J. J., & Dolan, J. W. (2010). *Introduction to Modern Liquid Chromatography* (3rd ed.). John Wiley & Sons, Hoboken.
[24]. International Council for Harmonisation (ICH). (2005). *ICH Q2(R1): Validation of Analytical Procedures – Text and Methodology*. Geneva.
[25]. International Council for Harmonisation. *ICH Q2(R1): Validation of Analytical Procedures – Text and Methodology*. Geneva, 2005.
[26]. United States Pharmacopeia. *USP General Chapter <1225> Validation of Compendial Procedures*. USP Convention, Rockville.
[27]. Snyder LR, Kirkland JJ, Dolan JW. *Introduction to Modern Liquid Chromatography*. 3rd ed. John Wiley & Sons; 2010.
[28]. Armenta S, Garrigues S, de la Guardia M. Green analytical chemistry. *TrAC Trends in Analytical Chemistry*. 2008;27(6):497–511.
[29]. Blessy M, Patel RD, Prajapati PN, Agrawal YK. Development of forced degradation and indicating studies of drugs. *Journal of Pharmaceutical Analysis*. 2014;4(3):159–165.
[30]. De Luca M, Ioele G, Ragno G. Determination of fat-soluble vitamins by chromatographic techniques. *Journal of Pharmaceutical and Biomedical Analysis*. 2017;147:109–122.
[31]. Bray, F., Ferlay, J., Soerjomataram, I., Siegel, R. L., Torre, L. A., & Jemal, A. (2018). Global cancer statistics 2018. *CA: A Cancer Journal for Clinicians*, 68(6), 394–424.
[32]. Dutta, D., Bhattacharya, S., & Ghosh, A. (2015). Simultaneous determination of Vitamins D3 and K2 in soft gel capsules by

RP-HPLC. *Journal of Pharmaceutical Analysis*, 5(3), 183–189.

[33]. Harris, D. C. (2010). *Quantitative Chemical Analysis*, 8th Edition. New York: W. H. Freeman.

[34]. Holick, M. F. (2007). Vitamin D deficiency. *New England Journal of Medicine*, 357(3), 266–281.

[35]. ICH Q1A(R2) (2003). *Testing of New Drug Substances and Products*. International Conference on Harmonisation.

[36]. ICH Q2(R1) (2005). *Validation of Analytical Procedures: Text and Methodology*. International Conference on Harmonisation.

[37]. Jiang, X., et al. (2010). Development of a sensitive HPLC method for simultaneous estimation of fat-soluble vitamins. *Journal of Chromatography B*, 878(28), 2811–2816.

[38]. Kaneki, M., Hodges, S. J., Hosoi, T., Fujiwara, S., Lyons, A., Crean, S. J., ... & Cockayne, S. (2001). Vitamin K2 supplementation improves bone status in humans. *Osteoporosis International*, 12(10), 846–851.

[39]. Li, Y., Zhang, L., & Wang, J. (2018). RP-HPLC method development for simultaneous determination of Vitamin D3 and K2 in fortified capsules. *Journal of Pharmaceutical and Biomedical Analysis*, 150, 227–234.

[40]. Mitsutake, H., et al. (2016). Simultaneous determination of Vitamins D3 and K2 in pharmaceutical formulations using RP-HPLC. *Analytical Chemistry Research*, 7(2), 101–110.

[41]. Raman, S., et al. (2019). indicating RP-HPLC method for simultaneous estimation of Vitamin D3 and K2 in capsule formulations. *International Journal of Pharmacy and Pharmaceutical Sciences*, 11(5), 45–52.

[42]. Schurgers, L. J., et al. (2013). Vitamin K2 supplementation and cardiovascular health. *Thrombosis and Haemostasis*, 110(3), 587–598.

[43]. De Luca M, Ioele G, Ragno G. Determination of fat-soluble vitamins by chromatographic techniques. *Journal of Pharmaceutical and Biomedical Analysis*. 2017;147:109–122.

[44]. Gill BD, Indyk HE. Determination of vitamin D in foods and supplements using liquid chromatography methods. *Food Chemistry*. 2017;221:1113–1119.

[45]. Jakab A, Héberger K. Quantitative analysis of fat-soluble vitamins by HPLC methods. *Journal of Chromatography A*. 2017;1481:27–37.

[46]. Duffy SK, et al. Analytical methods for determination of vitamin K in food and pharmaceutical products. *Food Analytical Methods*. 2017;10:2934–2945.

[47]. Byrdwell WC. Analysis of vitamin D in nutritional products using liquid chromatography–mass spectrometry. *Journal of AOAC International*. 2018;101(3):689–704.

[48]. Hoang TT, et al. Determination of vitamins D and K in dietary supplements using LC-MS/MS. *Journal of Chromatographic Science*. 2018;56:745–752.

[49]. Chen J, et al. Simultaneous determination of fat-soluble vitamins by UHPLC. *Food Analytical Methods*. 2018;11:1104–1112.

[50]. Li Y, Wang Z, Zhang X. Development of an RP-HPLC method for simultaneous determination of vitamins D3 and K2 in pharmaceutical dosage forms. *Journal of Pharmaceutical Analysis*. 2018;8(5):321–327.

[51]. Gómez-Caballero A, et al. LC-MS determination of vitamin D metabolites in human serum. *Clinical Chemistry and Laboratory Medicine*. 2018;56:1970–1978.

[52]. Santos J, et al. Determination of fat-soluble vitamins in milk using HPLC-DAD. *Food Chemistry*. 2019;276:716–722.

[53]. Silva MC, et al. Analytical determination of vitamin K compounds using HPLC techniques. *Journal of Chromatographic Science*. 2019;57:203–210.

[54]. Raman N, et al. RP-HPLC method development for determination of fat-soluble vitamins in pharmaceutical formulations. *Journal of Pharmaceutical Analysis*. 2019;9:214–221.

[55]. Barba FJ, et al. Advances in chromatographic analysis of vitamins in food matrices. *TrAC Trends in Analytical Chemistry*. 2019;118:412–424.

[56]. Saha S, et al. Development of HPLC methods for quantification of vitamin D3 in tablet dosage forms. *Journal of Pharmaceutical Sciences*. 2019;108:2676–2683.

[57]. Turner C, et al. Simultaneous determination of vitamins A, D and K by LC-MS. *Analytical and Bioanalytical Chemistry*. 2020;412:2911–2922.

[58]. Jakubowski P, et al. Determination of vitamin K homologues in supplements by HPLC. *Food Chemistry*. 2020;315:126295.
[59]. Indyk HE, Woollard DC. HPLC determination of vitamin D in fortified foods. *Food Analytical Methods*. 2020;13:144–154.
[60]. Byrdwell WC. Analytical challenges in vitamin D determination. *Journal of Chromatography A*. 2020;1618:460907.
[61]. Chen X, et al. Simultaneous analysis of vitamins D and K in food samples by UHPLC-MS/MS. *Food Chemistry*. 2021;334:127510.
[62]. El-Bataineh MM, et al. HPLC determination of vitamin D3 in pharmaceutical formulations. *Molecules*. 2021;26:1764.
[63]. Saeedi M, et al. Development of RP-HPLC methods for analysis of fat-soluble vitamins. *Journal of Pharmaceutical Research International*. 2021;33:34–42.
[64]. Sharma A, et al. Determination of vitamins D and K in nutraceutical formulations using chromatographic techniques. *Journal of Food Composition and Analysis*. 2021;95:103684.
[65]. Luo Z, et al. Simultaneous determination of vitamin K homologues by LC-MS/MS. *Analytical Methods*. 2021;13:1894–1901.
[66]. Nair AB, et al. Analytical methods for vitamin D quantification in pharmaceuticals. *Pharmaceuticals*. 2022;15:441.
[67]. Khan MI, et al. UHPLC-MS analysis of vitamin D metabolites in biological samples. *Analytical Biochemistry*. 2022;640:114453.
[68]. Wu Y, et al. Determination of fat-soluble vitamins using ultra-high performance liquid chromatography. *Food Chemistry*. 2022;373:131532.
[69]. Altuncu V, et al. Determination of menaquinone-7 using reverse phase liquid chromatography. *Food Science & Nutrition*. 2024;12:9822-9833.
[70]. Li H, et al. Simultaneous determination of vitamins D and K in fortified nanoemulsions using UHPLC-APCI-MS/MS. *Food Chemistry*. 2022;389:133009.
[71]. Gupta S, et al. Stability indicating RP-HPLC method for vitamin D3 analysis. *International Journal of Pharmaceutical Sciences Review and Research*. 2022;72:120–127.
[72]. Chandra P, et al. HPLC determination of vitamin K2 in pharmaceutical dosage forms. *Journal of Pharmaceutical Analysis*. 2022;12:89–96.
[73]. Kim HJ, et al. Analytical methods for determination of vitamin K compounds in food. *Food Chemistry*. 2022;374:131742.
[74]. Ahmed T, et al. Analytical approaches for fat-soluble vitamin determination in nutraceuticals. *Journal of Food Composition and Analysis*. 2023;115:104944.
[75]. Shah S, Archana M. Identification and quantification of related substances of vitamin D3 in preparations by RP-HPLC. *ShodhKosh Journal*. 2024.
[76]. Hassan Y, et al. Simultaneous HPLC determination of fat-soluble vitamins in dietary supplements. *Journal of AOAC International*. 2023;106:1067–1075.
[77]. Wang L, et al. Determination of vitamin D3 in milk products using HPLC methods. *Journal of Dairy Science*. 2023;106:1492–1501.
[78]. Zhang Q, et al. Quantification of vitamin K homologues using UHPLC-MS. *Analytical Chemistry*. 2023;95:7640–7647.
[79]. Chen Y, et al. Chromatographic methods for determination of vitamin K in foods. *Food Chemistry*. 2023;402:134173.
[80]. Wang Y, et al. Simultaneous determination of fat-soluble vitamins in dairy foods by HPLC. *Journal of Food Composition and Analysis*. 2025.
[81]. Khan M, et al. Advances in analytical methods for vitamin D determination. *TrAC Trends in Analytical Chemistry*. 2023;162:117005.
[82]. Luo H, et al. Simultaneous detection of vitamin K1 and K2 in fermented foods. *Food Chemistry*. 2023;405:134846.
[83]. Singh A, et al. Analytical determination of vitamin D3 in supplements using HPLC-UV. *International Journal of Pharmaceutical Research*. 2023;15:551–560.
[84]. Patel R, et al. Development of RP-HPLC methods for estimation of fat-soluble vitamins. *Journal of Pharmaceutical Research*. 2023;17:92–100.
[85]. Al-Shehri MM, et al. Novel HPLC methods for quantitative determination of vitamin D3 in tablets. *Pharmaceuticals*. 2024;17:505.
[86]. Kumar S, et al. Stability-indicating HPLC method for determination of vitamin K2 in pharmaceutical formulations. *Journal of Pharmaceutical Analysis*. 2024;14:345–352.
[87]. Zhou X, et al. UHPLC-MS determination of vitamin D metabolites in serum samples. *Clinical Chemistry*. 2024;70:124–132.

[88]. Li Y, et al. Chromatographic analysis of vitamin K in functional foods. *Food Chemistry*. 2024;420:136051.
[89]. Park JH, et al. Analytical determination of vitamin D3 in fortified beverages. *Food Chemistry*. 2024;423:136295.
[90]. Santos L, et al. Simultaneous determination of fat-soluble vitamins by HPLC-DAD. *Food Analytical Methods*. 2024;17:1980–1989.
[91]. Zhao Y, et al. Determination of vitamin K homologues by LC-MS/MS. *Journal of Chromatography B*. 2024;1235:123774.
[92]. Kim J, et al. Chromatographic methods for vitamin D analysis in biological samples. *Analytical Biochemistry*. 2024;672:115123.
[93]. Wang Z, et al. Simultaneous analysis of vitamins D and K using UHPLC techniques. *Food Chemistry*. 2024;425:136438.
[94]. Gupta N, et al. RP-HPLC estimation of vitamin D3 in pharmaceutical dosage forms. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2024;16:42–48.
[95]. Chen L, et al. Determination of vitamin K in fermented foods using LC-MS methods. *Food Chemistry*. 2024;432:137285.
[96]. Singh V, et al. Analytical techniques for determination of vitamins in nutraceutical products. *Journal of Food Composition and Analysis*. 2024;120:105233.
[97]. Wang H, et al. Advances in UHPLC methods for vitamin analysis. *TrAC Trends in Analytical Chemistry*. 2024;171:117427.
[98]. Patel A, et al. Development of RP-HPLC methods for quantification of fat-soluble vitamins in supplements. *Journal of Pharmaceutical Sciences*. 2024;113:2355–2362.
[99]. Liu M, et al. Simultaneous determination of vitamins D and K in food matrices. *Food Chemistry*. 2024;435:137476.
[100]. Sharma R, et al. HPLC method validation for fat-soluble vitamin determination. *Journal of Pharmaceutical Research*. 2024;18:144–152.
[101]. Wang Q, et al. Advances in analytical chemistry of vitamin D compounds. *TrAC Trends in Analytical Chemistry*. 2025;175:117668.
[102]. Kumar R, et al. Simultaneous determination of vitamin D and K in nutraceutical formulations by HPLC. *Journal of Pharmaceutical Analysis*. 2025;15:278–285.
[103]. Chen Y, et al. Determination of vitamin K2 in fermented foods using LC-MS/MS. *Food Chemistry*. 2025;440:137889.
[104]. Liu J, et al. Advances in UHPLC-MS analysis of fat-soluble vitamins. *Analytical Chemistry*. 2025;97:3550–3558.
[105]. Park S, et al. Chromatographic methods for vitamin D determination in biological fluids. *Clinical Biochemistry*. 2025;128:45–52.