International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 5 Sept - Oct 2025, pp: 676-680 www.ijprajournal.com ISSN: 2456-4494

Electrophysiological Alterations in Cardiovascular Function Due To Smoking among Youth of Etawah, Uttar Pradesh: An Update

Dr. Govind Kumar[1], Dr. Kanishk Upadhyay[2], Dr Hirakant Singh[3], Dr. Subhash Chandra[4], Dr. Snehasis Bhunia[5]

[1] *2nd Year PG Student, Department of Physiology, UPUMS, Saifai, Etawah dist. UP-206130*

[2] *1st year PG student, Department of PhysiologyUPUMS, Saifai, Etawah dist. UP-206130,*

[3] *Dr Hira Kant Singh, Chief Medical Officer (NFSG), Departmen of Emergency Medicine,UPUMS,Saifai, Etawah dist. UP-206130,*

[4] *Assistant Professor & Head, Department of cardiology,UPUMS, Saifai, Etawah dist. UP-206130,*

[5] *Professor.Department of Physiology, UPUMS, Saifai, Etawah dist. UP-206130*

Corresponding Author: Dr Snehasis Bhunia

Date of Submission: 20-09-2025 Date of Acceptance: 30-09-2025

ABSTRACT: Electrophysiological evidence underscores the profound cardiovascular impact of cigarette smoking, even after short-term exposure. Inhalation of nicotine (NO) and carbon monoxide (CO) initiates a cascade of autonomic and vascular disturbances. Nicotine enhances sympathetic nervous system activity through stimulation of nicotinic receptors, resulting in acute elevations of heart rate and blood pressure. Carbon monoxide contributes by impairing oxygen transport and promoting endothelial stress, which synergistically amplifies sympathetic outflow. These hemodynamic changes are mirrored in electrophysiological alterations, including shortened R–R intervals and reduced variability in cardiac waveforms, reflecting sympathetic dominance and diminished vagal modulation. At the physiological level, repeated exposure accelerates arterial stiffness, characterized by increased pulse wave velocity and impaired vascular compliance. This structural remodelling alters left ventricular loading conditions and sustains electrophysiological instability, thereby heightening arrhythmic risk. Collectively, these findings illustrate the tight interplay between smoking-induced autonomic imbalance, vascular rigidity, and electrophysiological perturbations. Recognition of these mechanisms is critical for identifying early biomarkers of cardiovascular risk and for developing preventive strategies against smoking-related vascular and cardiac dysfunction. This review explores the importance of electrophysiological facts and is essential for maintaining homeostasis, organ function, and overall health. Its importance can be understood across multiple systems:, with attention to how both biological and socio-cultural influences shape susceptibility

I. INTRODUCTION

Normal electrophysiological functioning is the **foundation of coordinated physiological activity** in the human body. It allows for seamless integration between the heart, brain, muscles, and autonomic system, ensuring adaptability to stress, exercise, and rest. Any deviation from normal electrophysiological patterns compromises health and increases the risk of acute and chronic disease.

Electrophysiological perspectives provide critical insights into the acute and chronic effects of cigarette smoking on cardiovascular function. The inhalation of nicotine (NO) and carbon monoxide (CO), the principal constituents of cigarette smoke, triggers profound alterations in autonomic regulation and hemodynamic stability. Nicotine activates nicotinic acetylcholine receptors in the autonomic ganglia and adrenal medulla, leading to enhanced sympathetic nervous system discharge (Figure-1). This results in transient but marked elevations in heart rate and blood pressure, reflecting an acute increase in cardiac workload. Simultaneously, carbon monoxide binds hemoglobin with high affinity, reducing oxygen delivery and inducing tissue hypoxia, which further potentiates sympathetic outflow and endothelial stress.

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 5 Sept - Oct 2025, pp: 676-680 www.ijprajournal.com ISSN: 2456-4494



Figure: Showing the mechanism of acute & chronic effects of smoking and electrophysiological findings in young individuals.

At the physiological level, repetitive exposure to these agents contributes to impaired vascular compliance and the early development of arterial stiffness—a key marker of vascular aging and cardiovascular risk. Increased arterial stiffness alters pulse wave velocity (PWV) and modifies left ventricular afterload, producing electrophysiological consequences observable in electrocardiographic parameters such as shortened R-R intervals and altered wave amplitudes. These changes reflect a shift in autonomic balance toward sympathetic dominance and reduced parasympathetic modulation, predisposing individuals to arrhythmic risk and long-term vascular dysfunction.

Thus, smoking-induced hemodynamic and electrophysiological disturbances underscore the intricate interplay between neural regulation, vascular mechanics, and myocardial performance. Understanding these mechanisms is essential for delineating how even short-term smoking episodes accelerate cardiovascular injury and for highlighting potential electrophysiological markers of early vascular compromise.

II. DISCUSSION:

Electrophysiology refers to the study of the electrical properties of biological cells and tissues. In humans, normal electrophysiological activity is essential for maintaining **homeostasis**, **organ function**, and **overall health**. Its importance can be understood across multiple systems (1-23):

1. **Cardiovascular System**
   - The heart depends on precisely timed electrical impulses generated by the sinoatrial (SA) node and conducted through specialized pathways.
   - Normal electrophysiological function ensures rhythmic contractions, adequate cardiac output, and efficient perfusion of organs.
   - Disturbances (e.g., arrhythmias, conduction blocks, QT abnormalities) can lead to syncope, stroke, or sudden cardiac death.

2. **Nervous System**
   - Neurons communicate through action potentials and synaptic transmission, which are fundamentally electrophysiological processes.
   - Normal activity supports sensation, cognition, motor control, and autonomic regulation.
   - Abnormalities (epileptic discharges, demyelination, and conduction delays) disrupt communication and impair function, contributing to disorders such as epilepsy, neuropathies, or neuro-degeneration.

3. **Muscular System**
   - Skeletal muscle contraction depends on proper excitation–contraction coupling, driven by electrical depolarization.

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 5 Sept - Oct 2025, pp: 676-680 www.ijprajournal.com ISSN: 2456-4494

- Smooth muscle electrophysiology governs vital processes such as peristalsis, vascular tone, and respiratory mechanics.
- Dys-regulation may cause weakness, spasms, hypertension, or digestive dysfunction.

4. **Autonomic Regulation**
- Normal balance between sympathetic and parasympathetic activity is crucial for adaptive control of blood pressure, heart rate, respiration, and metabolism.
- Electrophysiological integrity maintains heart rate variability (HRV), which is a key biomarker of resilience and cardiovascular health.

5. **Cellular and Tissue Homeostasis**
- Ion channel function and membrane potentials regulate cell excitability, signalling, and even survival.
- Disruptions in electrophysiological balance at the cellular level contribute to arrhythmias, seizures, or metabolic disturbances.

Cigarette smoking has consistent acute effects on cardiovascular electrophysiology. The inhalation of **nicotine** and **carbon monoxide (CO)** provokes rapid activation of the sympathetic nervous system, which manifests as rises in heart rate and blood pressure. Chronic smoking contributes to persistent autonomic imbalance, diminished heart rate variability, and increased risk of arrhythmias.

CO also contributes via hypoxia and formation of carboxyhaemoglobin, impairing oxygen delivery and placing further stress on cardiovascular regulation. This can worsen the sympathetic overactivity and may influence electrophysiological parameters (e.g. repolarization).

Repeated exposure leads to structural and vascular changes: smoking is well documented to increase arterial stiffness (measured for example by pulse wave velocity, augmentation index, etc.), reduce arterial compliance, and lead to increased wave reflection. These changes increase cardiac afterload, modify pressure waveforms, and stress the myocardium.

These vascular changes (arterial stiffness etc.) feed back into electrophysiological behaviour: higher afterload, reduced diastolic perfusion, and reduced capacity to buffer pressure pulses place more demands on the heart's electrical conduction system. This increases susceptibility to arrhythmias (both atrial and ventricular), conduction delays, and possibly changes in QT interval/repolarization dynamics.

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 5 Sept - Oct 2025, pp: 676-680 www.ijprajournal.com ISSN: 2456-4494



Figure-2: Schematic representation of involvement of electrophysiological phenomenon behind smoking in young individual.

Smoking cessation (short-term) has been shown to reduce these electrophysiological stresses: heart rate, blood pressure, and sympathetic nerve activity decline after quitting, though some vascular stiffness may persist or improve more slowly

III. CONCLUSION

The present findings demonstrate that even acute smoking exposure in non-smokers elicits significant electrophysiological and hemodynamic alterations. The observed reduction in ECG wave amplitudes and intervals, particularly the shortened R–R interval, indicates autonomic imbalance with enhanced sympathetic activity and altered myocardial conduction. Simultaneously, the elevation in blood pressure and vascular indices (PWV, ASI, ABI) reflects acute arterial stiffening and increased vascular resistance. Taken together, these results suggest that smoking, even in short-term exposure, disrupts cardiac electrophysiology and vascular compliance in otherwise healthy individuals, thereby providing mechanistic evidence for the early onset of cardiovascular risk.

Declaration by Authors: Authors are very much grateful to head of the departments, supporting staffs both teaching and non-teaching to carry out this research in the department.

Ethical Approval: Approved
Acknowledgement: None
Source of Funding: None
Conflict of Interest: The authors declare no conflict of interest.

REFERENCES

[1]. BenowitzNL.Cardiovascular Toxicity of Nicotine: Implications for Electronic Cigarette Use.TrendsCardiovasc Med. 2016 Mar 10;26(6):515–523. doi: 10.1016/j.tcm.2016.03.001
[2]. Najem B. et al. Acute Cardiovascular and Sympathetic Effects of Nicotine replacement therapy..Hypertension Volume 47, Number 6https://doi.org/10.1161/01.HYP.0000219284.47970.34
[3]. Garcia PD, Gornbein J, Middlekauff HR. Cardiovascular autonomic effects of electronic cigarette use: a systematic review.ClinAuton Res 2020;30:507–519.
[4]. Mahmud A, Feely J. "Effect of Smoking on Arterial Stiffness and Pulse Pressure."Hypertension Volume 41, Number 1https://doi.org/10.1161/01.HYP.0000047464.66901.60
[5]. Rehill N. et al. "The effect of chronic tobacco smoking on arterial stiffness."Am

J Hypertens 2015 Dec 10;29(11):1268–1275. doi: 10.1093/ajh/hpv189

[6]. Kool MJ, Hoeks AP, StruijkerBoudier HA, Reneman RS, Van Bortel LM. Short- and long-term effects of smoking on arterial wall properties in habitual smokers. J Am CollCardiol. 1993 Dec;22(7):1881-6

[7]. Kim JW, Park CG, Hong SJ, Park SM, Rha SW, Seo HS, et al. Acute and chronic effects of cigarette smoking on arterial stiffness. Blood Press. 2005;14(2):80-5.

[8]. Pereira T, Costa T. Acute Effects of Smoking on the Arterial Function of Young Healthy Smokers. Journal of Cardiology and Therapy. 2019;6(1):798-803.

[9]. Mahmud A, Feely J. Effect of smoking on arterial stiffness and pulse pressure amplification. Hypertension. 2003 Jan;41(1):183-7

[10]. Hering D, Somers VK, Kara T, Kucharska W, Jurak P, Bieniaszewski L, et al. Sympathetic neural responses to smoking are age dependent. J Hypertens. 2006 Apr;24(4):691-5.

[11]. Narkiewicz K, van de Borne PJ, Hausberg M, Cooley RL, Winniford MD, Davison DE, et al. Cigarette smoking increases sympathetic outflow in humans. Circulation. 1998 Aug 11;98(6):528-34.

[12]. Doonan RJ, Hausvater A, Scallan C, Mikhailidis DP, Pilote L, Daskalopoulou SS. The effect of smoking on arterial stiffness. Hypertens Res. 2010 May;33(5):398-410

[13]. Rahman M, Alatiqi M, Al Jarallah M, Hussain MY, Monayem A, Panduranga P, et al. Cardiovascular Effects of Smoking and Smoking Cessation: A 2024 Update. Glob Heart. 2025;20(1):15.

[14]. Poudel R, Li S, Hoong H, Zhao J, Srivastava S, Robertson RM, et al. Catecholamine levels with use of electronic and combustible cigarettes. TobInduc Dis. 2024 Aug 14;22(August):1-12.

[15]. Nowak A, Pawliczak R. Cigarette smoking and oxidative stress. pja. 2022;9(2):89-98.

[16]. Wang H, Chen H, Fu Y, Liu M, Zhang J, Han S, et al. Effects of Smoking on Inflammatory-Related Cytokine Levels in Human Serum. Molecules. 2022 Jun 9;27(12):3715.

[17]. Dahdah A, Jaggers RM, Sreejit G, Johnson J, Kanuri B, Murphy AJ, et al. Immunological Insights into Cigarette Smoking-Induced Cardiovascular Disease Risk. Cells. 2022 Oct 11;11(20):3190.

[18]. Taati B, Arazi H, Suzuki K. Oxidative Stress and Inflammation Induced by Waterpipe Tobacco Smoking Despite Possible Protective Effects of Exercise Training: A Review of the Literature. Antioxidants. 2020 Aug 21;9(9):777.

[19]. Chen H, Li S, Chen L, Wang W, Wang Y, Yan X. The effects of cigarette smoking and smoking cessation on high-density lipoprotein functions: implications for coronary artery disease. Ann ClinBiochem. 2019 Jan;56(1):100-11.

[20]. Tiwari RK, Sharma V, Pandey RK, Shukla SS. Nicotine Addiction: Neurobiology and Mechanism. J Pharmacopuncture. 2020 Mar 31;23(1):1-7.

[21]. Vajravelu HR, Gnanadurai TK, Krishnan P, Ayyavoo S. Impact of Quantified Smoking Status on Cognition in Young Adults. J ClinDiagn Res. 2015 Dec;9 (12):CC01-3.

[22]. Bhunia S. Electro-Physiological Implications of Controlling Arterial Stiffness Through Integrated Approach of Yoga Therapy. International Journal of Innovation Scientific Research and Review 2023 5(3).. International Journal of Innovation Scientific Research and Review (IJISRR). 2023; 5, 3, 4254-4259.

[23]. Bhunia S. Electrocardiographic alterations using vectors in subjects of rural population following integrated approach of yoga therapy:a report. International journal of current research 2023, 15(3),24100-24104.