International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 5 Sept - Oct 2025, pp: 1432-1439 www.ijprajournal.com ISSN: 2456-4494

Exploring the Synergistic Antioxidant and Antimicrobial Potential of Crassula ovata and Rosemary: A Phytopharmacological Perspective

Richa Singh[1] and Santosh Kumar Shukla[2]

1. Student, Department of Pharmaceutical Chemistry, Institute of Pharmaceutical Sciences & Research,Mahadev Campus, NH 25, Kanpur - Lucknow Rd, Sohramau, Lucknow, Lalpur, Uttar Pradesh 209859.

2. Assistant Professor, Department of Pharmaceutical Chemistry, Institute of Pharmaceutical Sciences & Research,Mahadev Campus, NH 25, Kanpur - Lucknow Rd, Sohramau, Lucknow, Lalpur, Uttar Pradesh 209859

Corresponding author:*Richa Singh*

Date of Submission: 01-11-2025 Date of Acceptance: 10-11-2025

ABSTRACT

Crassula ovata (Jade Plant) and Rosmarinus officinalis (Rosemary) are widely recognized medicinal plants traditionally valued for their healing, cleansing, and purifying properties. Both species are rich in diverse phytochemical constituents such as flavonoids, terpenoids, phenolic acids, and essential oils, which contribute to their broad pharmacological spectrum. The synergistic combination of these plants enhances their antioxidant and antimicrobial potential, as their bioactive compounds act complementarily to neutralize free radicals and inhibit microbial growth. This review focuses on the extraction techniques, phytochemical composition, and biological activities of C. ovata and R. officinalis, highlighting their promising roles as natural disinfectants, food preservatives, and herbal therapeutic agents. Collectively, these findings suggest that the integration of both plants could serve as a sustainable and effective approach in developing eco-friendly antimicrobial formulations and antioxidant-based health products.

Keywords: Crassula ovata, Rosmarinus officinalis, antioxidant activity, antimicrobial potential, phytochemistry, synergism, natural disinfectants.

I. INTRODUCTION

The rapid increase in microbial resistance and oxidative stress–related disorders has become a major global health concern, prompting researchers to explore safe, effective, and sustainable sources of bioactive compounds. Conventional synthetic antioxidants, antimicrobials, and disinfectants, though efficient, often cause undesirable side effects, cytotoxicity, or environmental hazards upon prolonged use. This growing awareness has shifted scientific interest toward natural plant-based compounds, which offer comparable efficacy with fewer toxic consequences. Medicinal plants have been traditionally used for centuries as reliable sources of therapeutic agents, owing to their rich phytochemical diversity and minimal adverse effects.

Among the many medicinally significant plants, Crassula ovata (commonly known as the Jade Plant) and Rosmarinus officinalis (popularly known as Rosemary) hold a prominent place in both traditional and modern herbal medicine. Crassula ovata, belonging to the family Crassulaceae, is a succulent plant widely cultivated for its ornamental as well as therapeutic value. It has been employed in traditional remedies for the treatment of wounds, skin infections, inflammation, and microbial ailments. Its fleshy leaves contain a variety of secondary metabolites such as flavonoids, triterpenes, tannins, alkaloids, and phenolics, which are known to contribute to its pharmacological effects. The plant exhibits significant antioxidant activity, which helps in neutralizing harmful free radicals and preventing oxidative cellular damage.

Rosmarinus officinalis, a member of the Lamiaceae family, is an aromatic evergreen shrub native to the Mediterranean region and now cultivated globally. It is widely recognized for its culinary, aromatic, and medicinal applications. The essential oil and extracts of rosemary are rich in bioactive compounds such as rosmarinic acid, carnosic acid, carnosol, camphor, and 1,8-cineole, which possess powerful antioxidant, antimicrobial, and anti-inflammatory properties. Traditionally, rosemary has been used as a natural antiseptic, preservative, and cognitive enhancer. Its antioxidant activity is largely attributed to phenolic diterpenes, which can scavenge reactive oxygen species and inhibit lipid peroxidation, thereby protecting biological systems from oxidative stress.

The combination of Crassula ovata and Rosmarinus officinalis presents a promising approach for enhancing biological efficacy through synergistic interactions. Synergy in phytotherapy occurs when the combination of bioactive compounds produces a more potent effect than the sum of their individual activities. In this case, the phenolic compounds and terpenoids from rosemary may enhance the solubility, stability, and absorption of flavonoids and triterpenes from C. ovata, leading to improved antioxidant and antimicrobial performance. Such a combination not only strengthens the pharmacological potential but also broadens the therapeutic spectrum against a range of oxidative and microbial challenges.

In addition to their therapeutic significance, these plants align well with the growing demand for natural and eco-friendly alternatives in pharmaceutical, cosmetic, and food industries. Extracts from C. ovata and R. officinalis can be incorporated into herbal disinfectants, wound-healing gels, oral care products, and antioxidant formulations as safe substitutes for synthetic agents. Their use in green formulations also supports environmental sustainability and reduces chemical load in ecosystems.

Therefore, the present study aims to explore the synergistic antioxidant and antimicrobial potential of Crassula ovata and Rosmarinus officinalis through a detailed analysis of their phytochemical composition, biological activities, and pharmacological applications. By understanding their individual and combined properties, this review seeks to provide a scientific basis for their integration into modern herbal formulations, offering a natural, safe, and effective approach to combating oxidative stress and microbial infections.

II. BOTANICAL DESCRIPTION

2.1. Crassula ovata (Jade Plant)

Crassula ovata, commonly known as the Jade Plant, belongs to the family Crassulaceae. It is a perennial, evergreen, succulent shrub characterized by its thick, fleshy, and glossy oval leaves that store water, an adaptation to survive in arid environments. The stems are stout and woody at the base, giving the plant a bonsai-like appearance, while the leaves are smooth, shiny, and dark green, sometimes edged with a reddish tint under bright sunlight. During the winter and early spring, the plant produces clusters of small, star-shaped pink to white flowers, which add to its ornamental appeal.

The Jade Plant is native to South Africa and Mozambique, thriving in dry, rocky regions, but it is now cultivated globally as an ornamental plant due to its low maintenance and adaptability. It is also commonly grown as a houseplant and is often associated with good fortune and prosperity in many cultures. Beyond its decorative use, C. ovata has notable ethnomedicinal importance. In traditional medicine, the leaf sap or paste is applied to treat wounds, cuts, burns, and warts, as it promotes healing and prevents microbial infections. Decoctions or extracts prepared from the leaves are used to alleviate inflammatory conditions, sore throats, and gastrointestinal disturbances. Scientific studies have confirmed that its leaves contain flavonoids, triterpenes, tannins, phenolic compounds, and alkaloids, which contribute to its antioxidant, anti-inflammatory, and antibacterial activities. Due to its wide range of bioactive constituents, C. ovata is now being explored as a promising plant for herbal drug formulation and natural therapy.

2.2. Rosmarinus officinalis (Rosemary)

Rosmarinus officinalis, commonly known as Rosemary, belongs to the family Lamiaceae. It is an evergreen, aromatic shrub that typically grows 1 to 2 meters in height. The plant is characterized by its slender, needle-like leaves, which are green on the upper surface and covered with fine, silvery hairs on the underside, giving them a grayish appearance. The plant produces small, bluish-purple flowers that bloom throughout the year, especially in warm climates, and its strong, refreshing aroma is due to the high concentration of essential oils present in its leaves.

Rosemary is native to the Mediterranean region but has now spread worldwide and is extensively cultivated for both culinary and medicinal purposes. It grows well in dry, sunny areas with well-drained soil and is often used as a garden herb for flavoring food. Beyond its culinary uses, R. officinalis holds a prominent place in traditional medicine and aromatherapy. Historically, it has been used as a memory enhancer, stimulant, and natural antiseptic. It is also employed in treating headaches, muscle pain, digestive problems, and respiratory ailments. The leaves and essential oil of rosemary are rich in phenolic diterpenes (such as carnosic acid and carnosol) and phenolic acids (like rosmarinic acid),

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 5 Sept - Oct 2025, pp: 1432-1439 www.ijprajournal.com ISSN: 2456-4494

which are known for their strong antioxidant and antimicrobial activities. Other essential oil constituents include camphor, 1,8-cineole, α-pinene, and borneol, which contribute to its aroma and pharmacological actions.

Due to these bioactive compounds, rosemary has been widely used in pharmaceutical, cosmetic, and food industries as a natural preservative and antioxidant agent. It not only prevents oxidative deterioration of food products but also protects biological systems from free radical–induced damage. Thus, R. officinalis serves as a multipurpose medicinal plant with remarkable therapeutic, nutritional, and aromatic value.

Table 1. Botanical Description of Crassula ovata and Rosmarinus officinali

Parameter	**Crassula ovata (Jade Plant)**	**Rosmarinus officinalis (Rosemary)**
Family	Crassulaceae	Lamiaceae
Common Name	Jade Plant, Money Plant	Rosemary
Plant Type	Succulent, perennial shrub	Evergreen aromatic shrub
Morphology	Fleshy, oval, glossy green leaves; woody stems; clusters of small pink-white, star-shaped flowers	Slender, needle-like leaves; bluish-purple flowers; strong aromatic odor
Height	1–1.5 meters	1–2 meters
Stem Characteristics	Thick, woody at base; smooth, greenish stems	Woody, erect stems with aromatic resin glands
Leaf Characteristics	Opposite, thick, fleshy, smooth leaves for water storage	Linear, narrow leaves with dark green upper surface and silvery underside
Flower Characteristics	Small, pinkish-white, star-shaped flowers in clusters	Small, blue to violet flowers in axillary clusters
Distribution	Native to South Africa and Mozambique; widely cultivated worldwide as ornamental and medicinal plant	Native to the Mediterranean region; cultivated globally for culinary, aromatic, and medicinal uses
Habitat	Prefers dry, rocky, and well-drained soil; drought-tolerant	Thrives in warm, dry climates with full sunlight and well-drained soil
Traditional Uses	Used for wound healing, inflammation, skin infections, sore throats, and microbial ailments	Used as memory enhancer, antiseptic, antioxidant, digestive aid, and natural preservative
Major Phytochemicals	Flavonoids, triterpenes, tannins, alkaloids, phenolics, ascorbic acid	Carnosic acid, carnosol, rosmarinic acid, camphor, 1,8-cineole, α-pinene, borneol
Pharmacological Activities	Antioxidant, anti-inflammatory, antimicrobial, wound healing	Antioxidant, antimicrobial, anti-inflammatory, neuroprotective, preservative
Economic Importance	Ornamental plant, herbal medicine, bioactive extract source	Culinary herb, cosmetic ingredient, herbal medicine, essential oil source

III. EXTRACTION METHODS

The extraction of bioactive compounds from medicinal plants plays a crucial role in determining their pharmacological efficacy. In the present study, the leaves of Crassula ovata and Rosmarinus officinalis are subjected to solvent extraction using methanol, ethanol, and aqueous solvents, as these solvents efficiently dissolve both polar and non-polar phytoconstituents. The Soxhlet extraction and maceration techniques are commonly employed to obtain concentrated crude extracts.

In the Soxhlet extraction method, the powdered plant material is continuously refluxed with a selected solvent for several hours, ensuring complete extraction of soluble bioactive compounds. Conversely, maceration involves soaking the powdered material in the solvent at room temperature for a specific duration, followed by filtration and concentration under reduced pressure.

For Rosmarinus officinalis, hydro-distillation is often utilized for the isolation of its essential oil, which is particularly rich in volatile monoterpenes such as 1,8-cineole, camphor, α-pinene, and borneol. These components contribute significantly to its antimicrobial and aromatic properties.

The extraction yield and phytochemical concentration are largely influenced by parameters such as solvent polarity, temperature, extraction duration, and particle size of the plant material. Methanol and ethanol typically yield extracts rich in phenolic compounds and flavonoids, while aqueous extracts are comparatively mild but safer for biological studies.

To evaluate potential synergistic bioactivities, combined extracts of Crassula ovata and Rosmarinus officinalis may be prepared in specific ratios such as 1:1 or 2:1. Such formulations allow investigation of possible interactions between their phytochemical constituents that could enhance antioxidant and antimicrobial properties.

IV. PHYTOCHEMICAL CONSTITUENTS

4.1. Crassula ovata (Jade Plant)

Crassula ovata, belonging to the family Crassulaceae, contains a wide range of secondary metabolites that contribute to its medicinal value. Phytochemical screening has revealed the presence of flavonoids, tannins, alkaloids, saponins, phenolic compounds, triterpenes, and ascorbic acid.

Among these, quercetin and kaempferol are the major flavonoids responsible for the plant's antioxidant and anti-inflammatory effects through free radical scavenging and enzyme inhibition mechanisms. Lupeol and β-sitosterol, which are triterpenoid compounds, play a vital role in membrane stabilization, wound healing, and antimicrobial defense. The presence of ascorbic acid (vitamin C) further enhances the antioxidant capacity of the plant, protecting cellular structures from oxidative damage.

Overall, Crassula ovata exhibits significant antioxidant and antibacterial potential, supporting its traditional use in treating wounds, skin infections, and inflammatory conditions.

4.2. Rosmarinus officinalis (Rosemary)

Rosmarinus officinalis, a member of the Lamiaceae family, is a well-documented medicinal herb rich in phenolic diterpenes and essential oils. The primary phenolic constituents include carnosic acid, carnosol, and rosmarinic acid, which possess potent antioxidant and neuroprotective properties.

In addition, rosemary's essential oil fraction contains volatile terpenoids such as 1,8-cineole, camphor, α-pinene, and borneol. These compounds are known for their antimicrobial, anti-inflammatory, and preservative properties. Phenolic acids in rosemary stabilize lipid membranes and inhibit oxidative degradation, thereby protecting biological systems from free radical-induced damage.

Due to this diverse phytochemical profile, R. officinalis has been extensively used in traditional medicine, aromatherapy, and the food industry as a natural antioxidant and antimicrobial agent.

4.3. Synergistic Phytochemical Interaction

The combination of Crassula ovata and Rosmarinus officinalis extracts exhibits synergistic effects, enhancing their overall therapeutic potential. Phenolic compounds such as rosmarinic acid and carnosic acid from rosemary can improve the solubility, stability, and bioavailability of flavonoids like quercetin and kaempferol from Crassula ovata.

This synergy results in stronger free radical scavenging, lipid peroxidation inhibition, and enhanced antimicrobial action compared to individual extracts. The complementary mechanisms of action—such as cell wall disruption by terpenoids and enzyme inhibition by flavonoids—contribute to broad-spectrum antibacterial activity.

Furthermore, the combination may reduce the effective concentration required for biological efficacy, minimizing potential toxicity and improving pharmacological safety. Thus, the synergistic interaction between these two botanicals supports their potential use as natural disinfectants, preservatives, and therapeutic agents in modern phytomedicine.

V. ANTIOXIDANT ACTIVITY

The increasing concern over oxidative stress and its involvement in degenerative diseases such as cancer, cardiovascular disorders, and neurodegeneration has intensified interest in natural antioxidants. Both Crassula ovata and Rosmarinus officinalis have demonstrated significant antioxidant potential due to their abundance of phenolic compounds, flavonoids, and terpenoids.

To evaluate the antioxidant potential, multiple in vitro assays such as DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging, ABTS (2,2′-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid) assay, FRAP (Ferric Reducing Antioxidant Power), and Hydrogen Peroxide (H_2O_2) scavenging assays are employed. These methods assess the capacity of plant extracts to neutralize free radicals or reduce oxidized intermediates.

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 5 Sept - Oct 2025, pp: 1432-1439 www.ijprajournal.com ISSN: 2456-4494

The methanolic and ethanolic extracts of Crassula ovata exhibit a dose-dependent radical scavenging activity, indicating that higher concentrations provide stronger inhibition of free radicals. This antioxidant behavior is primarily attributed to its flavonoid (quercetin, kaempferol) and triterpenoid (lupeol, β-sitosterol) content, which act as hydrogen or electron donors to quench reactive oxygen species (ROS).

Rosmarinus officinalis, on the other hand, demonstrates exceptionally high total phenolic content, with major contributors such as carnosic acid, carnosol, and rosmarinic acid. These compounds show strong correlation with total antioxidant capacity, often surpassing standard synthetic antioxidants like BHT or ascorbic acid in comparative studies.

When combined extracts of C. ovata and R. officinalis are tested, they display enhanced DPPH and ABTS radical scavenging activity, suggesting a synergistic interaction between phenolic diterpenes of rosemary and flavonoids of jade plant. This cooperative effect enhances redox potential and overall antioxidant efficiency.

The underlying mechanisms of antioxidant action include:

- **Donation of hydrogen atoms or electrons** to neutralize free radicals,
- **Chelation of pro-oxidant metal ions** (e.g., Fe^{2+}, Cu^{2+}) preventing Fenton reactions, and
- **Inhibition of lipid peroxidation**, thus protecting biological membranes from oxidative damage.

These findings highlight that the synergistic combination of both plant extracts could serve as a potent source of natural antioxidants for pharmaceutical, nutraceutical, and cosmetic applications.

VI. ANTIMICROBIAL ACTIVITY

Microbial infections caused by antibiotic-resistant strains have become a major global health concern, necessitating the search for novel plant-derived antimicrobial agents. Both Crassula ovata and Rosmarinus officinalis have demonstrated broad-spectrum antimicrobial activities, making them promising candidates for natural antimicrobial formulations.

The antimicrobial efficacy is generally evaluated using agar well diffusion and minimum inhibitory concentration (MIC) assays against a panel of pathogenic bacterial strains, including Escherichia coli, Staphylococcus aureus, Pseudomonas aeruginosa, and Bacillus subtilis.

Extracts of Crassula ovata exhibit significant inhibition zones against both Gram-positive and Gram-negative bacteria, attributed mainly to its triterpenoids, flavonoids, and phenolic acids. These bioactives disrupt bacterial cell walls, interfere with enzyme systems, and alter membrane permeability, leading to growth inhibition.

Rosmarinus officinalis essential oil demonstrates strong bactericidal action due to the presence of volatile monoterpenes such as 1,8-cineole, camphor, α-pinene, and borneol. These compounds penetrate microbial membranes, causing cytoplasmic leakage, protein denaturation, and enzyme inactivation. Additionally, rosmarinic and carnosic acids contribute to antimicrobial activity through oxidative stress modulation within microbial cells.

When both extracts are combined, they exhibit synergistic antibacterial effects, resulting in larger inhibition zones and lower MIC values compared to individual treatments. The synergism likely arises from complementary mechanisms, where rosemary terpenes enhance the permeability of microbial membranes, allowing the flavonoids and phenolics from C. ovata to act more effectively within the cell.

Furthermore, the combined extracts also show antifungal activity, notably against Candida albicans, where terpenoid–flavonoid interactions contribute to disruption of fungal cell walls and inhibition of ergosterol synthesis.

Overall, these findings indicate that the synergistic formulation of Crassula ovata and Rosmarinus officinalis can serve as a potent natural antimicrobial agent, suitable for applications in herbal disinfectants, topical formulations, and food preservation.

VII. MECHANISM OF ANTIMICROBIAL AND ANTIOXIDANT ACTION

The biological efficacy of Crassula ovata and Rosmarinus officinalis primarily depends on their rich phytochemical composition, which enables them to act through multiple biochemical mechanisms. Their antioxidant and antimicrobial effects are attributed to synergistic interactions between flavonoids, terpenoids, phenolic acids, and essential oils, which collectively contribute to cellular protection and pathogen inhibition.

7.1 Antioxidant Mechanisms

Both plants play a vital role in combating oxidative stress by scavenging reactive oxygen species (ROS) such as superoxide anions, hydroxyl radicals, and hydrogen peroxide, as well as reactive nitrogen species (RNS) including nitric oxide and peroxynitrite. These free radicals, when uncontrolled, lead to lipid peroxidation, DNA damage, and protein oxidation. The polyphenolic compounds—particularly quercetin, kaempferol, carnosic acid, and rosmarinicacid—act as hydrogen or electron donors, neutralizing these radicals and stabilizing them into non-reactive forms.

Additionally, these bioactives upregulate endogenous antioxidant defense enzymes such as superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx), thereby enhancing the cell's natural ability to counter oxidative stress. Some terpenoids also exhibit metal-chelating properties, preventing Fenton-type reactions that generate harmful hydroxyl radicals. Thus, the combined extract exerts a dual protective effect—direct scavenging of free radicals and stimulation of internal antioxidant systems.

7.2 Antimicrobial Mechanisms

The antimicrobial action of C. ovata and R. officinalis is multifaceted and targets different structural and metabolic pathways of microorganisms. One of the principal mechanisms involves the disruption of microbial cell membrane integrity. The lipophilic monoterpenes from rosemary (such as 1,8-cineole and camphor) integrate into bacterial membranes, causing increased permeability, ion leakage, and eventual cell lysis. Meanwhile, the flavonoids and triterpenoids from C. ovata inhibit bacterial enzymes involved in energy metabolism and protein synthesis, further impairing cellular function.

Another significant mechanism is the induction of oxidative stress within microbial cells, where the plant compounds promote the formation of intracellular ROS, leading to protein oxidation, nucleic acid degradation, and membrane damage. The synergistic phytochemical interaction between the phenolic compounds of rosemary and the flavonoids of jade plant enhances these bactericidal and fungicidal effects, resulting in broader antimicrobial coverage and higher potency than either extract alone.

VIII. PHARMACOLOGICAL AND INDUSTRIAL RELEVANCE

The pharmacological relevance of the synergistic extracts of Crassula ovata and Rosmarinus officinalis lies in their broad-spectrum antimicrobial and potent antioxidant activities, which make them valuable for multiple therapeutic and industrial applications.

In the pharmaceutical domain, these extracts can be incorporated into herbal disinfectants, topical wound-healing gels, and mouth rinses, serving as natural alternatives to chemical antiseptics. Their anti-inflammatory and tissue-regenerative properties further enhance their suitability for dermatological and oral care formulations.

In the cosmetic industry, the antioxidant-rich components protect the skin from oxidative damage and premature aging, making them ideal ingredients in anti-aging creams, serums, and lotions. Similarly, their antimicrobial potential helps in developing natural preservatives for cosmetic and personal care products, replacing synthetic parabens and phenolic compounds.

Moreover, the eco-friendly and biodegradable nature of these plant extracts positions them as safe alternatives to conventional sanitizers and preservatives that often cause environmental toxicity. Advanced applications involve nanoparticle-based delivery systems (liposomes, polymeric nanoparticles, and nanoemulsions) to improve bioavailability, stability, and controlled release of active constituents.

Furthermore, the synergistic antioxidant property of these plants opens potential for their inclusion in nutraceuticals and functional food products, offering preventive benefits against oxidative stress-related disorders such as cardiovascular diseases, diabetes, and neurodegeneration.

IX. FUTURE PROSPECTS

Although preliminary findings demonstrate promising antioxidant and antimicrobial efficacy, further research is necessary to fully explore and validate the therapeutic potential of Crassula ovata and Rosmarinus officinalis extracts.

Future studies should involve in vivo and clinical trials to confirm safety, bioavailability, and pharmacokinetics. Isolation of pure bioactive compounds through bioassay-guided fractionation can aid in identifying the most potent molecules responsible for synergistic effects.

Development of novel nanoformulations, such as liposomes, polymeric nanoparticles, or solid lipid nanoparticles, will enable targeted drug delivery and improved therapeutic efficiency. Evaluation of synergistic indices like the Fractional Inhibitory Concentration (FIC) will help quantify the degree of synergism and optimize dosage combinations for therapeutic applications.

Additionally, establishing standardization protocols for extract preparation—including solvent selection, extraction conditions, and quality control—will be essential to ensure reproducibility and pharmaceutical-grade consistency. Integrating these plant-based extracts into industrial applications can significantly advance the field of natural therapeutics and green formulation science.

X. CONCLUSION

The synergistic combination of Crassula ovata and Rosmarinus officinalis exhibits enhanced antioxidant and antimicrobial potential, attributed to the interactive effects of flavonoids, phenolic acids, and terpenoids. These phytochemicals function collectively to neutralize oxidative stress and inhibit a broad range of pathogenic microorganisms.

The dual pharmacological action supports their application as natural, safe, and eco-friendly alternatives to synthetic antioxidants and antimicrobial agents. Integration of these extracts into herbal disinfectants, cosmetic formulations, and nutraceuticals offers a sustainable approach to modern health and wellness products.

Future investigations focusing on mechanistic elucidation, clinical validation, and formulation optimization will be crucial to unlock their full therapeutic potential and facilitate their transition from traditional medicine to standardized pharmaceutical use.

REFERENCES

[1]. Ahmad, M., Khan, M. A., Zafar, M., & Sultana, S. (2014). Ethnobotanical uses and phytochemical screening of Crassula ovata in traditional medicine. Journal of Medicinal Plants Research, 8(15), 616–622.

[2]. Al-Sereiti, M. R., Abu-Amer, K. M., & Sen, P. (1999). Pharmacology of rosemary (Rosmarinus officinalis Linn.) and its therapeutic potentials. Indian Journal of Experimental Biology, 37(2), 124–130.

[3]. Amakura, Y., Umino, Y., Tsuji, S., &Tonogai, Y. (2000). Influence of rosmarinic acid and caffeic acid on oxidative stress in biological systems. Food and Chemical Toxicology, 38(2–3), 173–178.

[4]. Andrade, J. M., Faustino, C., Garcia, C., Ladeiras, D., Reis, C. P., & Rijo, P. (2018). Rosmarinus officinalis L.: An update review of its phytochemistry and biological activity. Future Science OA, 4(4), FSO283.

[5]. Aumeeruddy-Elalfi, Z., Gurib-Fakim, A., &Mahomoodally, M. F. (2016). Antimicrobial, antibiotic potentiating activity and phytochemical profile of essential oils from exotic and endemic medicinal plants of Mauritius. Industrial Crops and Products, 86, 111–119.

[6]. Basak, S., & Guha, P. (2018). Antioxidant and antibacterial activities of rosemary extracts in food preservation. Food Bioscience, 22, 87–94.

[7]. Benincasa, M., & Pizzo, F. (2017). The synergistic effect of plant phenolics in combating oxidative stress and microbial infections. Phytomedicine, 25, 75–84.

[8]. Blois, M. S. (1958). Antioxidant determinations by the use of a stable free radical. Nature, 181, 1199–1200.

[9]. Bozin, B., Mimica-Dukic, N., Simin, N., & Anackov, G. (2007). Characterization of the volatile composition of essential oils of Rosmarinus officinalis and their antioxidant and antimicrobial activities. Food Chemistry, 102(2), 302–308.

[10]. Brighente, I. M. C., Dias, M., Verdi, L. G., &Pizzolatti, M. G. (2007). Antioxidant activity and total phenolic content of some Brazilian species. Pharmaceutical Biology, 45(2), 156–161.

[11]. Burt, S. (2004). Essential oils: Their antibacterial properties and potential applications in foods—A review. International Journal of Food Microbiology, 94(3), 223–253.

[12]. Cao, G., & Prior, R. L. (1999). Measurement of oxygen radical absorbance capacity in biological samples. Methods in Enzymology, 299, 50–62.

[13]. Chen, Y., Zhang, H., & Liu, Z. (2020). Comparative antioxidant and antibacterial evaluation of rosemary extracts obtained by different extraction methods. Food Chemistry, 332, 127372.

[14]. Chiu, H. F., Venkatakrishnan, K., & Wang, C. K. (2018). Nutraceutical potential of rosemary and its constituents. Journal of Traditional and Complementary Medicine, 8(3), 383–390.
[15]. Dorman, H. J. D., & Deans, S. G. (2000). Antimicrobial agents from plants: Antibacterial activity of plant volatile oils. Journal of Applied Microbiology, 88(2), 308–316.
[16]. Edeoga, H. O., Okwu, D. E., &Mbaebie, B. O. (2005). Phytochemical constituents of some Nigerian medicinal plants. African Journal of Biotechnology, 4(7), 685–688.
[17]. Farhat, A., Jordán, M. J., Chaouech-Hamada, R., Landoulsi, A., & Sotomayor, J. A. (2017). Variations in essential oil, phenolic compounds, and antioxidant activity of Rosmarinus officinalis L. grown in different locations. Industrial Crops and Products, 107, 1–10.
[18]. Gachkar, L., Yadegari, D., Rezaei, M. B., Taghizadeh, M., & Rasooli, I. (2007). Chemical and biological characteristics of essential oil from Rosmarinus officinalis L. Food Chemistry, 102(3), 899–904.
[19]. Ghasemzadeh, A., & Ghasemzadeh, N. (2011). Flavonoids and phenolic acids: Role and biochemical activity in plants and human health. International Journal of Medical Sciences, 8(9), 843–858.
[20]. Halliwell, B., & Gutteridge, J. M. C. (2015). Free radicals in biology and medicine (5th ed.). Oxford University Press.
[21]. Inouye, S., Takizawa, T., & Yamaguchi, H. (2001). Antibacterial activity of essential oils and their major constituents against respiratory tract pathogens. Lett Appl Microbiol., 32(6), 481–485.
[22]. Kim, M. H., & Kim, Y. J. (2017). Synergistic antioxidant and antimicrobial potential of combined plant extracts. Pharmaceutical Biology, 55(1), 246–254.
[23]. Lu, Y., & Foo, L. Y. (2002). Polyphenolics of Rosmarinus officinalis and their antioxidant activities. Food Chemistry, 79(2), 221–230.
[24]. Maqsood, S., &Benjakul, S. (2010). Comparative antioxidant activity of aqueous and ethanolic extracts from leaves of selected herbs. Food Science and Technology International, 16(5), 461–468.
[25]. Mishra, B. B., & Tiwari, V. K. (2011). Natural products: An evolving role in future drug discovery. European Journal of Medicinal Chemistry, 46(10), 4769–4807.
[26]. Nazzaro, F., Fratianni, F., Coppola, R., & De Feo, V. (2017). Essential oils and antifungal activity. Pharmaceuticals, 10(4), 86.
[27]. Orhan, I. E., Özçelik, B., Kartal, M., & Sener, B. (2012). Antimicrobial and antiviral effects of some flavonoids and phenolic acids: A structure–activity relationship study. Chemistry & Biodiversity, 9(3), 485–495.
[28]. Parekh, J., Jadeja, D., & Chanda, S. (2005). Efficacy of aqueous and methanol extracts of some medicinal plants for potential antibacterial activity. Turkish Journal of Biology, 29(4), 203–210.
[29]. Singh, G., Maurya, S., Catalan, C., & De Lampasona, M. P. (2007). Chemical constituents, antimicrobial investigations and antioxidative potential of volatile oil of Rosmarinus officinalis. Food and Chemical Toxicology, 45(2), 232–240.
[30]. Singleton, V. L., Orthofer, R., &Lamuela-Raventós, R. M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by Folin–Ciocalteu reagent. Methods in Enzymology, 299, 152–178.