

Formulation and Assessment of the Antimicrobial Properties of Herbal Soap Made with Bioactive Neem Plant *Azadirachta Indica*

*Momoh Shaibu, Francisca O. Orivri

Department of Science Laboratory Technology
Federal Polytechnic Kaura Namoda, Zamfara State, Nigeria

Date of Submission: 15-08-2025

Date of Acceptance: 25-08-2025

ABSTRACT

In this research work, *Azadirachta indica* leaf extract was used to produce herbal soap as an eco-friendly product. The neem leaf samples were obtained from the Federal Polytechnic, Kaura Namoda campus. Ethanolic extraction process and concentration of crude extract were carried out. Different concentration of extract was used for the production of herbal soap: 10%, 20% and 30%. Phytochemical determination was carried out on the crude extract, and the following phytochemical compounds were determined: Alkaloid, Tannin, Saponins, terpenoid, and Phenol. Physicochemical assessment and antimicrobial activities were also carried out on the herbal soap produced. Moisture content, Total fatty matter, free alkali, free fatty acid, chloride content, pH, Form height, and hardness. All the values obtained fall within Iso/AOAC standard value for the toilet soap. The antimicrobial activities were examined against some clinical isolates of pathogenic microorganisms (*Staphylococcus aureus* and *Escherichia coli*) using the agar diffusion method. The pattern of inhibition varied with the soap concentration. The inhibition zone of each herbal soap against the test organism is *E. coli*: 1.45cm, 1.46 cm, and 1.65 cm for the concentrations of crude extract 10%, 20% and 30% respectively. While the inhibition zone against *Staphylococcus aureus*, 1.45cm, 1.46cm, and 1.55cm against the different concentrations of the extract. Therefore, the 30% crude extract used for the production of the herbal soap shows a high sensitivity against *E. coli*.

Keywords: Herbal Soap, *Azadirachta indica*, Phytochemicals, *Staphylococcus aureus*, *E. Coli*

I. INTRODUCTION

Skin infections are mainly caused by bacteria and fungi; the common bacteria are *Staphylococcus aureus* and *Streptococcus* species [1]. These infections are commonly treated with the aid of antibacterial drugs, creams, and anti-inflammatory agents. Other infections, such as

eczema, ringworm, and pruritus, are also treated with some succulent gel. Some used chemical substances, such as surfactants, to clean and wash the skin to prevent it from being contaminated by microorganisms [2].

Traditionally, people in rural areas often use various medicinal leaves to cure and prevent skin infections. The method and mode of using these medicinal leaves by villagers are boiling and bath with the extract for the skin infection. *Azadirachta indica*, commonly called the neem tree or neem leaves, appears green as a tropical tree that found almost everywhere in the northern part of Nigeria. The leaves and the stems of the neem tree are used for different medicinal purposes for both children and adults [3].

Several researchers have reported the antimicrobial effect of this neem extract, such as antidermatitic, anti-inflammatory, and other body protective properties. However, the tree is now commonly used for skin infection diseases such as staph and *E. coli* because of the bioactive phytochemicals present in the neem [4].

Therefore, innovating the neem extract to produce called Herbal soap as a vehicle to drive the bioactive compound into the skin tissue for effective use. These soaps are full of natural ingredients, such as essential oils and herbal extracts. The name herbal soap has become popularly accepted as medicinal soap for various skin infections. People love them because of their health benefits, they're environmentally friendly, and they don't contain all the harsh chemicals that regular soap does [5].

Many herbal soaps in the market today have a long history of their efficacy toward various skin diseases, most especially pathogens such as fungal and *Staphylococcus aureus* and other streptococcus species [6].

The literature has shown that most herbal soaps formulated by different researchers globally, the neem tree was part of the active ingredient for antimicrobial activities [7], [1],[4],[8], [9]. Researchers combine these leaf extracts with other

proven medicinal plants such as turmeric and Aloe vera extract.

Mechanisms of Action of Herbal Soap

Herbal soaps exert their various mechanisms of action against microorganisms. It penetrates the cell membranes because some of the compounds found in the herbal ingredient are integrated into the microbial cell membrane, therefore, causing the permeability and leakage of cell content to cell death. Herbal soap bioactive compounds can also inhibit the activity of enzymes such as protease and lipase and stop the replication of the microorganism [10].

It was reported that the bioactive compound in the herbal soap interferes with biofilm and making the microorganism more susceptible to eradication. Some herbal bioactive compound generates oxidative stress induction with the microbial cell wall. This action damages the protein, and lipids lead to the death of the cell [9].

Several studies have evaluated the efficacy of herbal soaps in clinical and laboratory settings. For instance, a study demonstrated a significant reduction in bacterial load on the skin compared to conventional soaps. Another study found that soap containing tea tree oil effectively reduced the presence of acne-causing bacteria, outperforming standard antimicrobial soaps. Comparative antimicrobial activities of Neem oil and its leaf extract have limited information in the literature [11].

Different extracting processes, such as the solvent used, may have also improved the activities of the bioactive compound of these neem extracts or juices [12]. Prepared a herbal using different plant extracts and found it effective on E. coli. The physiological understanding of the structure and basic functions of the skin, combined with literature evidence of natural and herbal remedies for common skin conditions as shown the importance of herbal soap. The benefit of herbal soap was not recognized until the global research findings showed the activities against various skin diseases [12]. Therefore, ethanol extract of neem leaves and the neem leaves powder were used to produce herbal soap using the cold saponification method, and the antimicrobial activities were tested and compared with isolated Staphylococcus and E. coli.

The growing interest in herbal soaps is driven by their perceived safety, natural origin, and potential health benefits. However, despite the popularity and purported antimicrobial properties of herbal soaps, there remain several critical

research gaps and challenges. This research would provide information that would be useful in the overview of herbal soap production with various bioactive plants [12].h

II. MATERIAL AND METHODS

All the chemicals and reagents used as purchased, they are of analytical grade. The equipment and materials used were sterilized, calibrated, and kept in a sterilized environment. The apparatus was selected and washed with detergents and rinsed with running tap water, and cleaned with distilled water.

SAMPLE COLLECTION

The neem leaves were collected from the Federal Polytechnic Kaura Namoda main campus into a sterilized polytene bag.

Extraction of bioactive compounds from neem leaves

The fresh neem leaves were cleaned with distilled water and spread in the laboratory for shade drying at room temperature. This process was chosen to prevent heat-sensitive compounds from damage at high temperatures. After 5 days, the dried material was ground to a fine powder with the aid of a laboratory mortar and pestle.

Cold extraction (Maceration) was used as described by 100 g of the dried neem leaves was weighed and 70% ethanol was also added in a sealed 1000 ml capacity conical flask. This was kept at room temperature for 72 hours with regular shaking. The solution was then filtered using Whatman filter paper number 1 to obtain the cleared filtrate. The filtrate was concentrated to remove the excess ethanol using a water bath at low temperature to evaporate the ethanol, and the semi-solid crude extract was used as a herbal ingredient for the soap production.

Soap Production using Cold Processes

The cold process of making soap is a common method for soap-making techniques. It is simply the process of mixing fat/oils with a strong alkali, such as sodium hydroxide, at room temperature, called saponification. An appropriate weight of sodium hydroxide was weighed and dissolved in distilled water, and kept for 24 hours. after that, the hydrometer gauge was used to determine the recommended value for soap production. The concentration of neem extract added varies in different concentrations in each soap production, ranging from 30%, 40% and 50%

neem extract. The herbal soap was kept for 5 days weeks in a dried place to allowed the harden.

Physicochemical Assessment of Herbal Soap

The physicochemical assessment was carried out on the herbal soap produced, weigh of the herbal soap, form height, form retention, and pH. The weight of the herbal soap was determined using a digital weigh balance to determine the actual weight of each herbal soap produced.

Form height was determined by weighing 1g of the herbal soap sample dissolved in 30ml of distilled water in a cleaned, calibrated measuring cylinder. 50 ml of distilled water was added and allowed to stand still until the aqueous layer was measured on each measuring cylinder and recorded. Form retention was also carried out by preparing 25 ml of 1% herbal soap samples into a cleaned measuring cylinder and shaking for 5 times, every one of each shake, the height was measured for about five times in one-minute intervals.

The pH test was carried out using a pH meter. 0.5 g of each herbal soap was dissolved in distilled water, the digital pH meter electrode was immersed in the soap solution, and the reading was taken for each soap solution.

ANTIMICROBIAL ACTIVITIES OF THE HERBAL SOAP

The test organisms used for this analysis were clinically isolated *Staphylococcus aureus*, *Escherichia coli*. The test organisms were prepared by streaking on a freshly prepared nutrient agar plate to obtain discrete colonies. A colony was picked with the aid of sterile wire loop and transferred into bottle containing sterile normal saline. the turbidity was compared with that of a McFarland turbidity standard scale 0.5 which is equivalent to a bacterial cell density of 1.5×10^8 CFU/ML [10].

Determination of Inhibitory Activity of The Soaps Using Agar Well Diffusion Method

Freshly prepared agar (20 ml) was prepared according to the method described by Aliyu et al. (2012, poured into sterile Petri-dishes, and the agar was allowed to solidify (gel). The bacterial strains were swabbed on the Nutrient agar plates. Four wells, each of diameter 6 mm, were made into each agar plate using a cork borer, and the plates were labeled. Unto each plate, 0.2 ml of appropriate soap dilution was placed in appropriate wells, that is, 300, 200, 100, and 50 mg/ml, respectively. The plates were left for 1 hour for the

herbal soap to diffuse into the agar and then incubated at 37 °C for 24 hours. After incubation, the plates were observed for evidence of inhibition (zone of inhibition). The diameters of the wells were measured using a calibrated ruler in cm. The experiment was performed in duplicates, and the averages were calculated.

III. RESULT AND DISCUSSION

The susceptibility of the two microorganisms (*S. aureus* and *E. coli*) to various concentrations of neem extract used for of the herbal soaps is shown in Table 1

Table 1: The result of inhibition of herbal soap with 10% neem extract on the bacterial

Test Organism	10% Neem extract of herbal soap	20% Neem extract of herbal soap	30% Neem extract of herbal soap
<u>E. Coli</u>	1.45cm	1.46cm	1.65cm
<u>S. aureus</u>	1.45cm	1.55cm	1.58cm

Key: E. Coli , Escherichia coli S. aureus = Staphylococcus aureus

Table 1 above shows the inhibition zone measured in (cm) against 10%, 20% and 30% concentration of neem extract used for the preparation of the herbal soap. The test organisms used were E. coli and S. aureus; the inhibition zone was measured as shown in the table above.

The 10% extract concentration has an inhibition zone of 1.45 cm on E. coli and S. aureus. The 20% neem extract showed a high inhibition zone on S. aureus (1.55 cm) and (1.46cm) on E. coli. The 30% neem extract of herbal soap has a high inhibition zone on E. coli (1.65 cm) and on S. aureus (1.58 cm). The statistical tool shows a significant difference among the different concentrations of Neem extract used in the preparation of the herbal soap. The antimicrobial activities of Neem extract have proven its efficacy on the E. coli and S. aureus, as it was also reported [7] that a herbal soap was prepared combination of different extracts, such as Neem extract and lemon grass[1], also reported the efficacy of Neem leaves extract on some bacterial species. Azadirachta Indica is grown in all parts of the world because of its medicinal value, as proven by different researchers [9], [13]. The antibacterial activities of Neem herbal soap produced for Neem oil show effectiveness on Staphylococcus aureus and Escherichia coli.

[13], developed Neem herbal soap with the combination of Tulsi powder. In their report, the herbal soap was proven effective on dermatophyte.

[14] Also prepared herbal soap using Azadirachta Indica and Ocimum sanctum, they concluded in their report that the combination has potential for antibacterial and antioxidant. [15] formulate and evaluate the neem herbal soap in India. They reported the action of producing herbal soap as very effective on anti-inflammatory, antihyperglycemic, antibacterial, and antifungal.

In the findings by the global researchers, Azadirachta Indica has proven its antimicrobial activities to be very effective on different kinds of skin diseases. The only variation on each finding by the different researchers is the method of

extraction of these Neem leaves, the concentration extract used, and the path of the Neem tree used to prepare herbal soap.




Figure 1. Culture Plate

Table 2. Quantitative Determination of Phytochemical Test on the Crude Neem Extract

Phytochemical	Test	Observation	Inference
Alkaloid	Dragendorff	Orange brown	Positive
Flavonoids	Shinoda test	No visible change	Negative
Phenol	Ferric Chloride test	Deep blue coloration	Positive
Glycoside	Keller-killiani test	No visible change	Negative
Terpenoid	Salkowski test	Golden yellowish	Positive
Tannins	Ferric chloride test	Blue-black coloration	Positive

From Table 2 above, a quantitative test for the presence of phytochemical compounds present from the neem leaves extract used for the formulation of herbal soap. The quantitative test shows the presence of Alkaloid, phenol, terpenoid, and tannins, while flavonoid and glucoside were absent. Phytochemicals terms as natural chemical compounds that are mostly found in plants. They have antimicrobial activities as reported by [16]. The antimicrobial activities of alkaloids, as reported by [7], are that alkaloid can interfere with and stop DNA replication or inhibit the essential enzymes of the microorganism because alkaloids contain nitrogen compounds. Polyphone compound has also been termed as a phytochemical that contains hydroxide groups; the activities of this compound interfere with the cell membranes of the microorganism because it causes the denaturation of microbial cell protein. On the other hand, the presence of terpenoids in the crude extract of the neem leaves shows its ability to penetrate the lipid bilayers of the microorganism, which in turn increases the permeability of the cell. The action of terpenoids as also been reported by [11] to it can cause oxidative stress and inhibit the enzyme responsible for microbial respiration. Finally, Tannin was quantitatively present in the crude Neem leaf extract. This also proves the evidence of herbal soap antimicrobial activities, because tannin

binds to the protein cell wall of microorganisms and result in inactive cell adhesion [17].

Table 3: Physicochemical Evaluation of Herbal Soap Produced

Parameter	Method	H.S1	H.S2	H.S3	ISO/AOAC value	STD
M. C %	Oven dried (105)	12%	12%	12%	10- 15 %	
TFM %	Titrimetric	78%	75,8%	74,7%	76 %	
F.A %	Titrimetric	0.21%	0.22%	0.18%	0.1 %	
F.F.A %	Titrimetric	1.6%	1.56%	1.48%	0.5 – 2.0%	
Chloride content %	Mohrs method	0.8%	0.8%	0.78	1.0%	
pH	pH meter	10.5	10.7	10.8	9-11	
F.H	Shake method (cylinder)	130mm	135 mm	140mm	>150 MM	
Hardness	Penetration test	Not crumble	Not crumble	Not crumbl e	-	
Insoluble alcohol matter (%)	Extraction method (using ethanol)	0.21%	0.34%	0.56%	<2.0%	

Key: M.C, moisture content, TFM, Total fatty matter, F.A, Free Alkali, FFA, Free Fatty Acid, F.H, Form height, H.S1, Herbal soap

Table 3 shows the physicochemical evaluation of the herbal soap produced from the neem leaf extract with different concentrations of the extract, 10%, 20% and 30% as shown in Table 3. The following parameters were determined, namely: moisture content, total fatty matter, free fatty acid, chloride content, pH, form height, and hardness. The moisture content of the herbal soap produced falls within the standard value of 10- 15% moisture content. The moisture content of soap determines the stability and shelf life.

The total fatty matter, free alkali content, and free fatty acid of the herbal soap produced fell within the standard range value of 76% for toilet soap, 0.1 % and 0.5- 2.0% respectively. However, the high total fatty matter determines the lather of the soap, while high free alkali may cause irritation of the skin. The chloride content, pH, form height, and hardness of the herbal soap produced are within the standard range; however, the chloride content of the soap produced is lower than the standard value, while the pH of the herbal soap falls between 10.5 -10.8, which is also within the range of standard value of 9- 11. This is to ensure that soap is safe to use for the skin [17].

IV. CONCLUSION

The research evaluates the antibacterial activities of herbal soap produced from different concentrations of the crude. The phytochemical compounds present are alkaloids, Tannin, Saponins, terpenoids, and Phenol. The Physicochemical assessment was also carried out on the herbal soap produced; Moisture content,

Total fatty matter, free alkali, free fatty acid, chloride content, pH, Form height, and hardness. A sensitivity assay was carried out on the herbal soap samples to determine the zone of inhibition. The results show that the pattern of inhibition varies with the soap concentration. The inhibition zone of each herbal soap against the test organism is *E. coli*: 1.45cm, 1.46 cm and 1.65 cm for the concentrations of crude extract 10%, 20% and 30% respectively. While the inhibition zone against *Staphylococcus aureus*, 1.45cm, 1.46cm, and 1.55cm against the different concentrations of the extract. Therefore, the 30% crude extract used for the production of the herbal soap shows a high sensitivity against *E. coli*

Conflict of interest: The authors declared no conflict of interest

Acknowledgment: The Authors acknowledge the Tertiary Education Trust Fund (TETFUND) through the Management of Federal Polytechnic Kaura Namoda, Zamfara State, Nigeria for the Sponsorship of this Research institutional base research (IBR)

REFERENCES

- [1]. N. Pooja, P. Mounika, D. Shailaja, N. Rakesh, J. Nandini, D. Keerthi Priya, R. Ramya, Potina Mounika (2025). Herbal Soap Formulation and Testing for Antibacterial and Skin Lightening Properties. World Journal of Pharmaceutical and Medical Research. 11(7), 371-377

- [2]. Neha ugale, Madhuri Darade, Sneha Kedar (2023). Formulation and evaluation of antimicrobial herbal soap. International Journal of Pharmaceutical Research and Applications 8(2) DOI:10.35629/7781-080318441850
- [3]. Gupta Vijay Prabhu Mamgain Shilpa (2024). Formulation and Evaluation of Antibacterial Herbal Soap. International Journal of Creative Research Thoughts (IJCRT). 11(6) www.ijcrt.org
- [4]. Disha Arora¹, Asha Gandhi, Amanjeet Singh, Rohit Kumar (2024). Formulation and Evaluation of Antimicrobial Herbal Soap. International Journal of Research and Analytical Reviews (Ijrar) 11(2) www.ijrar.org
- [5]. John Lloyd Bernaldez¹ and Manuela Cecille G. Vicencio (2021). Antibacterial Activity of Soap Formulated from Garlic (*Allium sativum* L.) Extract. Journal of Advances in Microbiology 21(1): doi: 10.9734/JAMB/2021/v21i130320
- [6]. Warra AA, Wawata IG, Umar RA, Gunu, S.Y. (2012) Soxhlet extraction, Physicochemical Analysis and Cold process Saponification Nigeria *Jatropha curcas* L. seed oil. Canadian J.Pure and Appl. Sci. 6(1):1803-180719
- [7]. Ameh, A. O., Muhammad, J. A., and Audu, H. G. (2013). Synthesis and characterization of antiseptic soap from neem oil and shea butter oil. Africa Journal of Biotechnology 12(29), pp. 4656-4662 64
- [8]. A.Seetha Devi, D.V.Sivani, D.Anusha, G. Sarath, Syed Meraj Sultana (2021). Formulation aand Evaluation of Antimicrobial Herbal Soap. International Journal of Pharmaceutical Sciences Review and Research (71)2. <http://dx.doi.org/10.47583/ijpsrr.2021.v71i02.019>
- [9]. Sri DwiYanti, Siti Sulandjari, T., Winanti, I.,Asto L. A. (2021). CypirusRotundus L: Formulation and Evaluation Antiseptic Soap Advances in Engineering Research, volume 209 International Joint Conference on Science and Engineering (2021)
- [10]. Aliyu, M.S., Tijjani, M.B., Doko, M.H., Garba, M.M. Ibrahim, S.M. Abdulkadir, D., Abba U.U. (2012). Antimicrobial Activity of SabulunSalo a Local Traditional Medicated Soap Nigerian Journal of Basic and Applied Science (2012), 20(1): 35-38. <http://www.ajol.info/index.php/njbas/index>
- [11]. Wylie MR and Merrell DS (2022). TheAntimicrobial Potential of the NeemTree Azadirachtaindica.Front. Pharmacol. 13:891535.doi: 10.3389/fphar.2022.891535
- [12]. Afshan Ashraf and Khadija Rafiq (2025). Preparation and characterization of antibacterial herbal soap. Journal of Pharmacognosy and Phytochemistry 2025; 14(1): 388-397 <https://doi.org/10.22271/phyto.2025.v14.i1e.15261>
- [13]. Mahesh D. Shinde, Shravan J. Somani (2023). Development, Design and Evaluation of Herbal NeemSoap. International Journal of inPharmaceutical Sciences 1(12). Doi:10.5281/Zenodo.10431834
- [14]. Rajesh D Mokate, Jayashree S Kokat, Puja B Aher, Amruta Kakade, Shraddha Patil and Dnyaneshwari Darade (2024). Preparation and evaluation of herbal antimicrobial soap. Journal of Pharmacognosy and Phytochemistry 13(5)<https://doi.org/10.22271/phyto.2024.v13.i5d.15089>
- [15]. Sandip Tole, RutujaKhaire, SayyedMukhim Sayyed Salim,Gulshan Rathi (2024). Formulation aand Evaluation of Herbal Neem Soap. International Research Journal of Modernization in Engineering Technology and Science 6(6). www.irjmets.com
- [16]. Keshav S., Tejpal Y., Gaurav D., Ravindra P. (2023). Application of Phytoconstituents in Herbal Soap Formulation. J. Adv. Res. in Pharmaceutical Sciences and Pharmacology Interventions6(2)2. <https://orcid.org/0000-0001-7673-6127>
- [17]. Raja Kumar¹, Md Shoib Akhtar, Mansi Gupta (2023). Formulation and Evaluation of Herbal Neem andTurmeric SoapEpra International Journal Of Research And Development (Ijrd) 8(4), Doi:10.36713/Epra2016