International Journal of Pharmaceutical research and Applications
Volume 11, Issue 1, Jan - Feb 2026, pp:67-71 www.ijprajournal.com ISSN: 2456-4494

Nature's Healing Touch: A Review of Herbal Gels for the Treatment of Mouth Ulcers.

Pankaj Pappu Jadhav*, Devesh Pravinkumar Bhavsar, Dr. Ravindra S. Sonawane

*1. Student * KES's Late Shri P.C.B.College of D.Pharmacy &
Late Prof.R.K.kele College of B.Pharmacy, Amalner, Dist Jalgaon, India
2. Assistant Professor, KES's Late Shri P.C.B.College of D.Pharmacy &
Late Prof.R.K.kele College of B.Pharmacy, Amalner, Dist Jalgaon, India
3. Principal, KES's Late Shri P.C.B.College of D.Pharmacy &
Late Prof.R.K.kele College of B.Pharmacy, Amalner, Dist Jalgaon, India
*Corresponding Author

Date of Submission: 02-01-2026 Date of acceptance: 10-01-2026

ABSTRACT:-
Mouth ulcers are a common oral health concern that cause pain, discomfort, and impaired quality of life. Conventional treatments often provide symptomatic relief but may be associated with side effects or limited efficacy, prompting interest in natural alternatives. Herbal gels formulated with bioactive agents such as aloe vera, turmeric, tulsi, and honey have emerged as promising therapeutic options due to their anti-inflammatory, antimicrobial, antioxidant, and wound-healing properties. Aloe vera contributes mucoprotective and soothing effects, while turmeric offers curcumin-mediated anti-inflammatory and antimicrobial activity. Tulsi (Ocimum sanctum) enhances immune modulation and provides broad-spectrum antimicrobial benefits, and honey accelerates tissue regeneration through its osmotic, antioxidant, and antibacterial actions. Collectively, these natural agents demonstrate synergistic potential in reducing pain, promoting healing, and preventing secondary infections in oral ulcer management. This review critically evaluates the pharmacological basis, formulation strategies, and clinical evidence supporting herbal gels containing aloe vera, turmeric, tulsi, and honey, highlighting their role as safe, effective, and accessible alternatives for the treatment of mouth ulcers.

Keywords:- Mouth Ulcer, Natural Ingredients, wound healing agents, Mucoadhesive polymers

I. Introduction:-

This review aims to explore the scientific basis, formulation strategies, and clinical relevance of herbal gels containing aloe vera, turmeric, tulsi, and honey in the treatment of mouth ulcers. By consolidating evidence from traditional medicine and modern research, the article highlights the potential of these natural agents to serve as effective, accessible, and patient-friendly alternatives to conventional therapies.

In recent years, there has been growing interest in natural remedies and herbal formulations as safer, more holistic alternatives. Herbal gels, in particular, offer a convenient topical delivery system that ensures prolonged contact with the lesion site, enhancing therapeutic efficacy. Among the wide array of natural agents, aloe vera, turmeric, tulsi (Ocimum sanctum), and honey stand out due to their well-documented pharmacological properties. Aloe vera is valued for its soothing, anti-inflammatory, and wound-healing effects. Turmeric, rich in curcumin, exhibits potent anti-inflammatory and antimicrobial activity. Tulsi contributes immunomodulatory and antioxidant benefits, while honey provides antibacterial action and accelerates tissue repair through its osmotic and antioxidant mechanisms.

II. PATHOPHYSIOLOGY OF MOUTH ULCERS:-

The exact mechanism of mouth ulcer formation is not fully understood; however, immune-mediated tissue damage plays a key role. Triggering factors stimulate the release of pro-inflammatory cytokines such as tumor necrosis factor-alpha (TNF-α), leading to epithelial breakdown and ulcer formation. Oxidative stress and microbial colonization further delay healing. An ideal therapeutic agent should therefore reduce inflammation, control microbial growth, neutralize free radicals, and promote tissue regeneration.[3,6,7,9]

International Journal of Pharmaceutical research and Applications
Volume 11, Issue 1, Jan - Feb 2026, pp:67-71 www.ijprajournal.com ISSN: 2456-4494

III. RATIONALE FOR HERBAL MOUTH ULCER GELS:-

Herbal medicines have been traditionally used in oral care due to their biocompatibility and therapeutic versatility. Herbal mouth ulcer gels offer several advantages:

- Localized drug delivery at the site of action
- Reduced systemic absorption and toxicity
- Synergistic action of multiple phytoconstituents
- Better patient compliance due to soothing effect
- The gel dosage form is particularly suitable as it forms a protective layer over the ulcer, reduces irritation, and ensures sustained release of active constituents.[9,11,12,14]

IV. Natures touch herbal mouth ulcer gels:-

1. Aloe vera

Aloe vera gel contains polysaccharides, amino acids, vitamins, and enzymes that promote epithelial regeneration. It exhibits anti-inflammatory activity by inhibiting prostaglandin synthesis and accelerates wound contraction and healing.[11,12]

2. Curcuma longa (Turmeric)

Curcumin, the principal bioactive compound, possesses strong antioxidant and anti-inflammatory properties. It suppresses inflammatory mediators and enhances granulation tissue formation, aiding faster ulcer healing.[9,10]

3. Ocimum sanctum (Tulsi)

Tulsi exhibits antimicrobial, immunomodulatory, and analgesic activities. It is effective against oral pathogens and helps prevent secondary infections at ulcer sites.[13,14]

4. Centella Asiatica

This herb promotes collagen synthesis and angiogenesis, essential for tissue repair. Triterpenoids present in the plant accelerate wound healing and improve epithelialization.[15,16]

5. Honey and Propolis

Honey acts as a natural antimicrobial agent and maintains a moist wound environment. Propolis contains flavonoids that reduce inflammation and enhance tissue regeneration.

Other herbs such as chamomile, neem, clove, and licorice are also commonly incorporated for their soothing and antimicrobial effects.[17,18,19]

V. Organoleptic characteristics and evaluation tests of selected herbal ingredients:-

Ingredient	Physical Form / Order	Color	Odor	Taste	Organoleptic Properties	Tests Performed for Identification & Quality Evaluation
Aloe vera	Semi-solid gel / extract	Colorless to pale green	Mild, characteristic	Slightly bitter	Smooth, slippery texture with cooling sensation	pH test, viscosity determination, polysaccharide (acemannan) test, microbial limit test
Curcuma longa (Turmeric)	Fine powder	Bright yellow to orange	Strong, aromatic	Bitter and pungent	Dry, fine powder with intense yellow pigmentation	Curcumin identification test, alkali color reaction test, loss on drying, ash value
Ocimum sanctum (Tulsi)	Dried leaf powder / extract	Greenish-brown	Pleasant, aromatic	Pungent, slightly bitter	Coarse powder with strong herbal aroma	Volatile oil estimation, microscopic evaluation, extractive value test
Centellaasiatica	Powder / extract	Light green to brown	Mild, grassy	Slightly bitter	Soft texture with herbal odor	Triterpenoid confirmation test, total ash value, moisture content analysis
Honey	Viscous liquid	Golden yellow to amber	Pleasant, floral	Sweet	Thick, sticky, smooth consistency	Reducing sugar test, viscosity measurement, pH determination, adulteration test
Propolis	Resinous mass / extract	Dark brown	Strong, balsamic	Sharp, slightly bitter	Sticky, resin-like texture with intense aroma	Flavonoid identification test, solubility test, melting point determination

VI. Polymers and excipients used in formulation:-

1. Gelling Agents

- The choice of polymer significantly affects viscosity, spreadability, mucoadhesion, and drug release:
- Carbopol 934/940 – excellent mucoadhesive and viscosity properties
- Hydroxypropyl methylcellulose (HPMC) – non-irritant and film-forming
- Chitosan – natural polymer with antimicrobial activity
- Sodium alginate – provides smooth texture and bioadhesion. [20 to 26]

2. Other Excipients

- Humectants (glycerin, propylene glycol): prevent drying
- Preservatives (sodium benzoate, parabens): inhibit microbial growth

- Flavoring agents: improve palatability
- Sweeteners (xylitol): enhance taste and reduce cariogenic risk. [27,29]

VII. Formulation methodology of herbal mouth ulcer gel:-

1. Extraction of Herbal Actives-Herbal materials are shade dried, powdered, and extracted using aqueous or hydro-alcoholic solvents. Extraction methods include maceration, percolation, and Soxhlet extraction. The extract is filtered, concentrated, and stored under controlled conditions.[20,22,33]
2. Preparation of Gel Base-The selected polymer is dispersed in distilled water and allowed to hydrate completely. The pH is adjusted between 6.5 and 7.0 to match oral physiological conditions.[21,26]
3. Incorporation of Herbal Extract-The herbal extract is slowly added to the hydrated gel base with continuous stirring to ensure uniform distribution.[25,30]
4. Final Adjustment and Packaging-Excipients are added, and the formulation is evaluated for consistency, homogeneity, and stability before being packed in collapsible tubes or suitable containers.[27,28]

VIII. Evaluation parameters:-

1. Organoleptic Evaluation-Assessment of color, odor, texture, and homogeneity to ensure patient acceptability.
2. pH Measurement-Ensures compatibility with oral mucosa and prevents irritation.
3. Viscosity and Rheological Behavior-Determines ease of application and retention at the ulcer site.
4. Spreadability-Measures the ability of the gel to spread evenly over mucosal surfaces.
5. Drug Content Uniformity-Ensures uniform distribution of active constituents.
6 .Mucoadhesive Strength-Evaluates the gel's ability to adhere to oral mucosa for prolonged action.
7. In Vitro Drug Release-Determines release kinetics using diffusion cell methods.
8. Antimicrobial Activity-Assessed against oral pathogens such as Streptococcus mutans and Candida albicans.
9. Stability Studies-Conducted as per ICH guidelines to evaluate physical and chemical stability.[23,24,25,28,31,32,33,34,35]

IX. Conclusion:-

Mouth ulcers remain a prevalent oral health problem that significantly impacts daily life, and their effective management continues to be a clinical priority. While conventional therapies provide symptomatic relief, concerns regarding side effects and recurrence have encouraged exploration of safer, natural alternatives. Herbal gels incorporating with natural ingredients demonstrate promising therapeutic potential due to their combined anti-inflammatory, antimicrobial, antioxidant, and wound-healing properties. Aloe vera soothes and protects mucosal tissue, turmeric reduces inflammation and microbial load, tulsi enhances immune defense, and honey accelerates tissue repair while preventing secondary infection. The synergistic action of these agents not only alleviates pain and discomfort but also promotes faster healing, making herbal gels a viable adjunct or alternative to conventional treatment. Moreover, their accessibility, affordability, and cultural acceptance strengthen their relevance in both clinical and community settings. Future research should focus on standardized formulations, controlled clinical trials, and mechanistic studies to validate efficacy and ensure consistent therapeutic outcomes.

References:-

[1]. 1.Scully C, Porter S. Br J Oral Maxillofac Surg. 2008;46:198–206.
[2]. 2.Preeti L et al. J Oral MaxillofacPathol. 2011;15:252–256.
[3]. Ship JA et al. Quintessence Int. 2000;31:95–112.
[4]. Edgar NR et al. J ClinAesthetDermatol. 2017;10:26–36.
[5]. Scully C. N Engl J Med. 2006;355:165–172.
[6]. Altenburg A et al. DtschArztebl Int. 2014;111:665–673.
[7]. McBride DR. Am Fam Physician. 2000;62:149–154.
[8]. Barrons RW. Am J Health Syst Pharm. 2001;58:41–50.
[9]. Aggarwal BB, Harikumar KB. Int J Biochem Cell Biol. 2009;41:40–59.
[10]. Chainani-Wu N. J Altern Complement Med. 2003;9:161–168.
[11]. Surjushe A et al. Indian J Dermatol. 2008;53:163–166.
[12]. Chithra P et al. Mol Cell Biochem. 1998;181:71–76.
[13]. Cohen MM. J Ayurveda Integr Med. 2014;5:251–259.
[14]. Pattanayak P et al. Pharmacogn Rev. 2010;4:95–105.

[15]. Brinkhaus B et al. Phytomedicine. 2000;7:427–448.
[16]. Shukla A et al. Phytother Res. 1999;13:50–54.
[17]. Molan PC. J Wound Care. 1999;8:415–418.
[18]. Kujumgiev A et al. J Ethnopharmacol. 1999;64:235–240.
[19]. Cowan MM. ClinMicrobiol Rev. 1999;12:564–582.
[20]. Lachman L et al. Industrial Pharmacy. Lea &Febiger; 1986.
[21]. Aulton ME. Aulton's Pharmaceutics. Elsevier; 2018.
[22]. Gupta GD, Gaud RS. Indian J Pharm Sci. 2005;67:585–588.
[23]. Smart JD. Adv Drug Deliv Rev. 2005;57:1556–1568.
[24]. Andrews GP et al. Eur J Pharm Biopharm. 2009;71:505–518.
[25]. Patel RP et al. Int J Drug Deliv Technol. 2009;1:41–51.
[26]. Kaur LP, Guleri TK. J Drug DelivTher. 2013;3:1–6.
[27]. Indian Pharmacopoeia. IPC; 2022.
[28]. ICH Q1A(R2). Stability Testing Guidelines; 2003.
[29]. Jain A et al. Asian J Pharm Clin Res. 2014;7:51–55.
[30]. Prasanth VV et al. Int J Pharm Pharm Sci. 2011;3:120–123.
[31]. Dash GK, Murthy PN. Indian J Pharm Sci. 2011;73:371–374.
[32]. Sharma PK, Yadav S. Int J Pharm Sci Rev Res. 2015;32:58–65.
[33]. Gupta A et al. J Pharm SciInnov. 2012;1:45–50.
[34]. Panigrahi L et al. Indian J Pharm Sci. 2006;68:246–249.
[35]. Prabu GR et al. J ApplMicrobiol. 2006;101:487–495.