International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 290-298 www.ijprajournal.com ISSN: 2456-4494

Opinions on the Severity, Identification, and Appropriate Therapies for Tonsillitis (Inflammatory Infection of the Tonsils) in Recent Years, as well as Nowadays, Circumstances

Yash Srivastav[1*], Nutan Shrivastava[2], Madhaw Kumar[3], Nikita Sharma[4], Mohammad Aqil Siddiqui[5], Jyotsana Shrivastava[6]

[1]*Shri Venkateshwara University, Gajraula, Uttar Pradesh, India. (Assistant Professor)*
[2]*City Women's College, Jankipuram, Lucknow, Uttar Pradesh, India.*
[3]*Goel Institute of Pharmacy & Sciences, Lucknow, Uttar Pradesh, India.*
[4]*Signa College of Pharmacy, Kanpur, Uttar Pradesh, India.*
[5]*Lucknow College of Pharmacy,Lucknow, Uttar Pradesh, India.*
[6]*Dr. Rammanohar Lohia Avadh University, Uttar Pradesh, India.*

Date of Submission: 05-11-2025 Date of Acceptance: 15-11-2025

ABSTRACT

An inflammation of the tonsils in the upper throat is called tonsillitis. It may be acute or long-term. Tonsillitis that is acute usually develops quickly. A scratchy throat, fever, tonsil enlargement, difficulty swallowing, and swollen lymph nodes around the neck are possible symptoms. Peritonsillar abscess (quinsy) is one of the complications.Celsus wrote about tonsillectomy with sharp instruments in the first century AD, followed by vinegar and other medicinal rinses. Physicians have been recording tonsillitis treatments ever since. In the late 19th century, tonsillitis began to receive more medical attention.Known as Waldeyer's tonsillar ring, the tonsils (TON-sills) are a group of lymphoid organs that face the aerodigestive tract. They are made up of the lingual tonsils, two tubal tonsils, two palatine tonsils, and the adenoid tonsil, also known as the pharyngeal tonsil. These organs are crucial to the immune system.In contrast to younger children who are impacted by viral aetiology, children between the ages of 5 and 15 often acquire tonsillitis due to Streptococcus bacteria. According to reports, 17.82% of school-age children in India had chronic tonsillitis.Inflammation of the tonsils, or tonsillitis, accounts for about 0.4% of outpatient visits in the US. In 70% to 95% of cases, a viral infection is the cause of tonsillitis. However, 5% to 15% of adults and 15% to 30% of patients aged five to fifteen have tonsillitis due to bacterial infections caused by group A beta-hemolytic streptococcus (Streptococcus pyogenes). Because group A beta-hemolytic streptococcus infections can progress to more systemic complications like abscess, acute glomerulonephritis, rheumatic fever, and scarlet fever, it's critical to distinguish them from other bacterial or viral causes of pharyngitis and tonsillitis. There are numerous diagnostic methods available, such as oropharyngeal and serum laboratory testing, as well as symptom-based validated rating systems (such as the Centor score).Males were more frequently impacted. Due to awareness of the similarities between the symptoms of tonsillitis and COVID-19, there was a significant increase in the number of patients complaining of sore throats. Groups with lower socioeconomic status were particularly impacted. The two most prevalent forms of tonsillitis were follicular and parenchymatous.Tonsillitis can be treated at home by gargling with salt water, consuming warm beverages, eating cold, soft foods, and more. Here are some pain-relieving tips.The cause of tonsillitis determines the course of treatment. The goal of treating viral tonsillitis is to control symptoms. Antibiotics may be used to treat bacterial tonsillitis. All ages can get tonsillitis, although children between the ages of five and fifteen are most likely to get it.Maintaining proper hygiene is one of the best strategies to avoid this. The infections that cause tonsillitis are communicable, but the illness itself is not. One of the best methods to lower your risk of infection is to wash your hands frequently, stay away from others who have colds, and try not to touch your face, mouth, or nose when you're out and about. Warm beverages, cold foods, gargling salt water, and more are some at-home self-care remedies for tonsillitis. Tonsil irritation is referred to as tonsillitis. Two oval-shaped glands located at the rear of the throat are called tonsils. Their job is to combat germs and viruses that enter the body through the mouth and nose.It discusses tonsillitis's causes, diagnosis, course of therapy, and risks.

Keywords: Warts, Epidemiology, Aetiology, Treatments

I. INTRODUCTION

Tonsillitis is an inflammatory tonsil infection brought on by microorganisms—typically hemolytic streptococci bacteria or viruses—invading the mucous membrane. The symptoms include fever, lethargy, swallowing difficulties, sore throat, and swollen lymph nodes on both sides of the neck.A viral infection is typically the cause of tonsillitis, or inflammation of the tonsils, which accounts for about 0.4% of outpatient visits in the US. Droplets from patients with acute tonsillitis or, in rare cases, asymptomatic carriers are the means of transmission. Recurrent tonsillitis may be the cause of tonsilloliths, or tonsil stones, although the precise aetiology is unknown, and the calcified buildup of food particles, cell debris, and microbes in deep invaginations on the tonsils' surface results in tonsilloliths, also known as tonsillar stones. Tonsilloliths greater than 5 mm are uncommon, but tonsilloliths smaller than 5 mm are common. Although the precise process of tonsillolith development is unclear, recurrent tonsillitis may be a factor.A physical examination is typically the only method used to diagnose tonsilloliths. Computed tomography of the oropharynx can be useful if the diagnosis is unclear, especially if the stone has not extruded or is large. This would show a radiopaque, calcified, cylindrical or spherical lesion in the tonsillar tissue or fossa. Tonsilloliths can also be detected using panoramic radiography (1). Expectant management is used for tonsilloliths. Gargling with saltwater and brushing your teeth frequently can help treat halitosis. Surgery is rarely necessary if tonsilloliths grow too big to pass on their own. Tonsillolith-related complications are uncommon and are documented as case reports in the literature. Necrosis of the compressed tissue, as well as compression and displacement of surrounding tissues, can result from large tonsilloliths.About 1.3% of outpatient visits are caused by tonsillitis, a common inflammatory illness affecting the palatine tonsils. The illness is usually brought on by bacterial or viral infections, and symptoms include tonsillar erythema or exudate, fever, odynophagia, and sore throat. Although bacterial causes, particularly group A Streptococcus, can result in serious consequences such as peritonsillar abscess, severe rheumatic fever, and poststreptococcal glomerulonephritis, viral tonsillitis typically resolves on its own. To differentiate bacterial from viral etiologies, prompt and precise diagnosis is crucial. Microbiological testing and clinical scoring methods like the Centor or McIsaac criteria assist in direct, proper care and cut down on needless antibiotic use. To avoid complications and encourage a successful recovery, early detection and management are essential.Viral or bacterial infections are the main cause of tonsillitis, an inflammatory disease of the palatine tonsils that accounts for 1.3% of outpatient visits. Between the anterior palatoglossal arch and the posterior palatopharyngeal arch, in the lateral oropharynx, are the tonsils (2-4). In 70% to 95% of cases, a viral infection is the cause of tonsillitis. Adenovirus, coronavirus, respiratory syncytial virus, and rhinovirus are the most frequent viral culprits. Rubella, hepatitis A, CMV, and Epstein-Barr virus are less frequent causes. Group A beta-hemolytic streptococcus (GABHS) infections cause tonsillitis in 5% to 15% of adults and 15% to 30% of patients aged 5 to 15. Children under five years old are more likely to have viral aetiologies, and children under two years old are less likely to have GABHS.The tonsils function as immunological barriers as part of Waldeyer's ring, but inflammation of the tonsils can cause problems like rheumatic fever, post-streptococcal glomerulonephritis, and peritonsillar abscess. To avoid needless antibiotic use and reduce consequences, accurate diagnosis and treatment are essential.The lateral area of the oropharynx contains the palatine tonsils, also called the pharyngeal tonsils. Known as the palatine arches or pillars, they are situated between the anterior palatoglossal arch and the posterior palatopharyngeal arch. The tonsils, along with the adenoids (nasopharyngeal tonsil), tubal tonsil, and lingual tonsil, comprise Waldeyer's ring, a protective lymphatic arrangement. The tonsils are composed of lymphatic tissue. By capturing and filtering dangerous microbes, they serve as the body's first line of defence against infections that enter through the digestive or respiratory systems. They also play a crucial part in the immune response.The most common symptom of the illness, which is mostly brought on by bacterial or viral infections, is a sore throat. Clinical evaluation is the main method used to diagnose acute tonsillitis. It can be difficult to distinguish between bacterial and viral origins, yet doing so is crucial to preventing needless antibiotic use, highlighting the necessity of precise assessment to provide appropriate treatment.With a focus on current recommendations for antimicrobial therapy and criteria for surgical referral, this training activity offers evidence-based guidance on the diagnosis and management of tonsillitis. Students learn about the concepts of antimicrobial stewardship and approved diagnostic instruments (5-8). In order to improve diagnostic accuracy, optimise treatment regimens, and minimise needless interventions, the course emphasises the importance of interprofessional collaboration among doctors, nurses, pharmacists, and laboratory staff. By preventing overtreatment and preventing the emergence of antibiotic resistance, coordinated care improves patient-centred outcomes and lessens the strain on healthcare systems.Acute tonsillitis is usually diagnosed clinically and has an infectious cause. 70% to 95% of cases are caused by viral pathogens, but 5% to 15% of adult cases and 15% to 30% of pediatric infections, particularly in children aged 5 to 15, are caused by bacterial pathogens, primarily Streptococcus pyogenes (group A Streptococcus [GAS]). In order to minimise needless antibiotic prescriptions and promote effective antimicrobial use, it is crucial to promptly distinguish between bacterial and viral tonsillitis. This differentiation is important because consequences from streptococcal tonsillitis include acute rheumatic fever, poststreptococcal glomerulonephritis, and peritonsillar abscess (9).

Fig.1 Tonsillitis

EPIDEMIOLOGY

About 2% of ambulatory contacts in the US are caused by tonsillitis, which is a frequent reason for outpatient visits. Although it can occur at any time of year, it is most prevalent in winter and early spring. In some age groups, tonsillitis is mostly caused by Group A beta-hemolytic Streptococcus (GABHS), which accounts for 15% to 30% of infections in children aged 5 to 15 and 5% to 15% in adults. On the other hand, GABHS is rare in children under the age of two, while viral infections are more common in children under the age of five. Hospital admissions for severe or difficult tonsillitis have increased recently in some areas; this trend may be related to a decrease in tonsillectomy rates following the implementation of stricter surgical criteria. Significant morbidity and a lower quality of life are linked to recurrent tonsillitis, which is defined as five or more bouts per year.Age-group differences in the etiological distribution are substantial. 5–15% of adult instances of pharyngitis and 15–30% of cases are caused by Group A beta-hemolytic streptococcus (GABHS). in kids between the ages of 5 and 15, making it the most frequent bacterial cause in kids in school. On the other hand, viral infections are more common in children under five, and GABHS is especially uncommon in children under two. These epidemiological trends emphasise how crucial age-specific diagnostic methods are. Younger children usually have viral infections that can be treated without antibiotics, while teenagers and school-age children are more likely to need testing for GABHS (10-13). Clinicians can reduce needless antibiotic use and ensure proper care of bacterial cases by optimising diagnostic and treatment techniques with an understanding of these trends.Fever, sore throats, tonsillar exudates, and sensitive anterior cervical lymphadenopathy are characteristic symptoms of acute tonsillitis. Other symptoms include dysphagia (difficulty swallowing) and odynophagia (painful swallowing), which are frequently brought on by inflammation and swelling of the tonsils. Accurate diagnosis and treatment depend on a thorough history and physical examination. Clinicians should evaluate the patient's sexual history and vaccination status, especially for influenza and diphtheria, since tonsillar involvement can be a symptom of several infections, such as syphilis, HIV, and gonorrhea. 5,627 patients with tonsillitis or quinsy were admitted at that time; 2,376 of them were adults (42%) and 3,251 were children (58%). Student's t-test and linear regression were used to compute statistical probability. A tonsillectomy was performed on 203 children and 127 adults in 2003. Despite minimal increases in tonsillectomy rates, local tonsillitis admissions have grown by 118% in adults and 179% in children since 2003. Nationwide, comparable patterns have been noted. The results of this study could be seen as the outcome of the National Health Service's current policy. In a university teaching hospital, tonsillitis admissions and tonsillectomy rates were analysed retrospectively during a ten-year period from 2003 to 2012. Despite minimal increases in tonsillectomy rates, local tonsillitis admissions have grown by 118% in adults and 179% in children since 2003 (14-17).

ETIOLOGY

Infectious agents are the main cause of tonsillitis, with viral infections responsible for 70%–95% of cases. Rhinovirus, respiratory syncytial virus, adenovirus, coronavirus, Epstein-Barr virus (EBV), cytomegalovirus, and human immunodeficiency virus are examples of common viral etiologies. Although systemic viruses like EBV can cause more severe or protracted disease, these viruses typically appear as part of more general upper respiratory tract infections. Due to their potential for complications, bacterial causes of tonsillitis are clinically significant even if they are less common. The most common bacterial infection, especially in children between the ages of five and fifteen, is Group A beta-hemolytic Streptococcus, often known as S pyogenes. Staphylococcus aureus, Streptococcus pneumoniae, Haemophilus influenzae, Fusobacterium necrophorum (particularly in adolescents and young adults), and Streptococcus dysgalactiae are additional species implicated.Mycoplasma pneumoniae, Chlamydophila pneumoniae, and Neisseria gonorrhoeae are less prevalent bacterial infections, especially in sexually active people. In unvaccinated people, Corynebacterium diphtheriae should be taken into consideration. Chronic or recurring tonsillitis has also been linked to Mycobacterium tuberculosis, especially in high-risk groups. Biofilm-forming bacteria like S aureus and H influenzae contribute to the persistence of recurrent tonsillitis, which is frequently polymicrobial. Recurrence has also been linked to changes in the tonsillar microbiota and decreased microbial diversity (dysbiosis).Suppurative and nonsuppurative tonsillitis consequences include scarlet fever, rheumatic fever, acute glomerulonephritis, and abscess; however, these are uncommon in the developed world.

A deposit of pus between the tonsillar capsule and the pharyngeal constrictor muscle is known as a peritonsillar abscess. Fever, sore throat, dysphagia, trismus, and a "hot potato" voice are among the symptoms. A palatine bulge lateral to the tonsil may be visible to the doctor upon examination. The most frequent consequence of streptococcal tonsillitis is a peritonsillar abscess. Antibiotic treatment for tonsillitis decreases the development of abscesses, despite the fact that the two are clinically different (18-22).

DIAGNOSED

Although systematic grading systems and microbiological tests can help differentiate between bacterial and viral etiologies, tonsillitis is primarily diagnosed clinically. The Centor and FeverPAIN (fever in the last 24 hours, purulence, attend quickly [within 3 days], inflamed tonsils, and no cough or coryza) scores are examples of tools that enhance diagnostic precision and direct the need for additional testing or antibiotic medication.Microbiological Testing: Particularly in probable cases of group GABHS infection, microbiological confirmation aids in distinguishing between bacterial and viral tonsillitis. Preventing needless antibiotic usage and supporting targeted treatment are two benefits of accurately identifying the causal culprit. Quick test for antigen detection. In children, the rapid antigen detection test (RADT) has a high specificity (88%–100%) but a varied sensitivity (61%–95%). While a negative result in high-suspicion pediatric patients should prompt additional testing, a positive result confirms the diagnosis of streptococcal pharyngitis. Culture of the throat is regarded as the best standard for diagnosing GABHS, especially in kids with strong clinical suspicion but negative RADT results. Before starting antibiotics, cultures are necessary, according to the guidelines of the Infectious Diseases Society of America.Patient age, clinical appearance, and risk factors should all be taken into account when testing for tonsillitis. Due to the low risk of streptococcal sequelae, including rheumatic fever, routine testing is generally not advised for children under three. When infectious mononucleosis is suspected, monospot or EBV serology can be helpful (23-25). For sexually active people, a nucleic acid amplification test or culture for Neisseria gonorrhoeae or Chlamydia trachomatis should be taken into consideration. Patients with unusual symptoms or high-risk exposures may benefit from fast plasma reagin and human immunodeficiency virus testing.Laboratory and Imaging Studies: Although they are not usually necessary for simple tonsillitis, laboratory and imaging studies may be necessary in some situations to check for complications or other diagnosis. These tests can verify suspected pathogens, determine systemic involvement, or check for deep neck space infections. Based on the severity, duration, and related results, clinical judgment should direct their use. Laboratory tests: In cases of simple tonsillitis, routine blood work, such as a metabolic panel or complete blood count, is usually not required. However, when evaluating patients with dehydration or systemic toxicity, or when taking into account other diseases, these tests might be helpful. Imaging studies: When issues are suspected, imaging is used.The recommended method is neck computed tomography (CT), which has a high sensitivity for identifying deep space infections and defining the extent of abscesses. Ultrasound may occasionally be used as a first modality, particularly in children or in situations where CT is not easily accessible.Clinical Scoring Tools: Based on factors including tonsillar exudates, tender anterior cervical lymphadenopathy, fever, and lack of cough, the Centor score is a clinical tool used to estimate the likelihood of GAS pharyngitis. A higher score raises the likelihood of a bacterial infection and can influence choices about testing or empirical antibiotic treatment. Each positive result is worth one point (26-28).

TREATMENT OR MANAGEMENT

The underlying cause, intensity of symptoms, and likelihood of complications all influence how tonsillitis is managed. The majority of cases are self-limiting and viral, necessitating just supportive care. Antibiotic medication may be necessary for bacterial infections, especially those brought on by GAS, in order to shorten the length of symptoms and avoid consequences. Supportive Care: The majority of tonsillitis cases are viral, self-limiting, and only need symptomatic treatment. Adequate water, rest, and analgesics such as acetaminophen or nonsteroidal anti-inflammatory medications are advised interventions. Warm salt-water gargles and throat lozenges could offer more relief. There is still little evidence to justify the occasional use of topical medications such as viscous lidocaine. Zinc gluconate and herbal therapies are generally not advised due to their uneven benefits. Adjunctive Corticosteroids: In patients with severe odynophagia in particular, a single dosage of corticosteroids (such as dexamethasone) may hasten the remission of symptoms and offer temporary pain relief. Particularly in patients with diabetes or immunocompromised conditions, where possible hazards may outweigh the advantages, use should be customised. Antibiotic Therapy: GAS is the most prevalent cause of proven or highly suspected bacterial tonsillitis, for which antibiotics are recommended. Penicillin V or amoxicillin for ten days, azithromycin, clarithromycin, or first-generation cephalosporins for patients with nonanaphylactic penicillin allergy are examples of first-line treatments.Antibiotics may reduce the length of symptoms by about 16 hours, but their main advantage is that they prevent complications

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 290-298 www.ijprajournal.com ISSN: 2456-4494

like acute rheumatic fever and peritonsillar abscess, especially in high-risk groups (e.g., Indigenous Australians or underprivileged communities). Antibiotic resistance, gastrointestinal distress, Clostridium difficile infection, and higher medical expenses are possible dangers. Due to the low risk of transmission and complications, chronic GAS carriers usually do not need treatment (29). According to current guidelines from the American College of Physicians and the Centers for Disease Control and Prevention, antibiotic therapy is only advised when GAS is verified by throat culture or fast antigen detection tests. In low-risk situations, a cautious waiting strategy or delayed prescription might be suitable.Chronic versus Recurrent Tonsillitis: Chronic tonsillitis may be connected to Helicobacter pylori infection and is linked to a higher risk of immunoglobulin A nephropathy, which may have wider implications for both local and systemic health. These extra-tonsillar consequences may be caused by immunological activation and persistent inflammation in chronic tonsillitis. A common definition of recurrent tonsillitis is five or more occurrences per year. Assessment for underlying disorders like immunodeficiency should be part of the evaluation. The following are guidelines for medical therapy: In recurring cases, clindamycin or amoxicillin-clavulanate may be more effective than penicillin in eliminating GAS. Low-dose azithromycin or long-acting benzathine penicillin are examples of prophylactic antibiotics that can lower the frequency of recurrences. Because of its superior adherence, azithromycin is frequently chosen for youngsters.Surgical Management: When a patient's quality of life is greatly impacted by the frequency or severity of recurring or chronic tonsillitis episodes, tonsillectomy is recommended. When a patient is not responding to medical treatment or has issues such a recurrent peritonsillar abscess, this surgery may also be considered. Children with recurrent GAS-related throat infections may be eligible for tonsillectomy according to the Paradise criteria, which are evidence-based recommendations (30-33).



Fig. 2Management of Tonsillitis

II. DISCUSSION AND CONCLUSION

Our review articles address tonsillitis's epidemiology, diagnostics, alternative treatments, and several causes in addition to providing a broad summary of the condition. Despite the fact that tonsillitis can be treated with herbal-related medications and other therapies, our results suggest that further clinical study is necessary because tonsillitis is a complicated condition. More randomised controlled studies are required to treat tonsillitis. In the future, we hope to offer a preliminary tonsillitis assessment. Future counselling-based research in our state or country will assess patients' mental and physical health with the assistance of our colleagues in order to deliver

International Journal of Pharmaceutical Research and Applications
Volume 10, Issue 6 Nov - Dec 2025, pp: 290-298 www.ijprajournal.com ISSN: 2456-4494

more accurate information about tonsillitis and its treatment.

ETHICAL STATEMENT

A pharmacist needs to act with integrity and honesty. A pharmacist refrains from actions that undermine their dedication to acting in their patients' best interests, such as discriminatory acts, inappropriate behaviour, and unfavourable working conditions that cloud judgment. A pharmacist keeps up their level of expertise.

ACKNOWLEDGEMENT

The authors would like to thank Shri Venkateshwara University, Gajraula, Uttar Pradesh, India, City Women's College, Jankipuram, Lucknow, Uttar Pradesh, India,Goel Institute of Pharmacy & Sciences, Lucknow, U.P., India & Signa College of Pharmacy, Kanpur, Uttar Pradesh, India, for extending their facilities.

CONFLICT OF INTEREST

The authors attest that they are free of any known financial or personal conflicts of interest that would taint the findings of this study.

INFORMED CONSENT

Using websites, review articles, and other sources to produce research content.

REFERENCES

[1]. Meegalla N, Downs BW. StatPearls [Internet]. StatPearls Publishing; Treasure Island (FL): Jun 5, 2023. Anatomy, Head and Neck, Palatine Tonsil (Faucial Tonsils).

[2]. Anderson J, Paterek E. Tonsillitis. StatPearls. April 30, 2022. Updated September 18, 2022. Accessed January 31, 2022. https://www.ncbi. nlm.nih.gov/books/NBK544342/

[3]. Windfuhr JP, Toepfner N, Steffen G, et al. Clinical practice guideline: tonsillitis I. Diagnostics and nonsurgical management. Eur Arch Otorhinolaryngol. 2016;273(4):973-987.

[4]. Georgalas CC, Tolley NS, Narula A. Tonsillitis. BMJ Clin Evid. 2014;2014: 0503.

[5]. Bartlett A, Bola S, Williams R. Acute tonsillitis and its complications: an overview. J R Nav Med Serv. 2015;101(1):69-73. [PubMed]

[6]. Smith KL, Hughes R, Myrex P. Tonsillitis and Tonsilloliths: Diagnosis and Management. Am Fam Physician. 2023 Jan;107(1):35-41. [PubMed]

[7]. Putukian M, McGrew CA, Benjamin HJ, Hammell MK, Hwang CE, Ray JW, Statuta SM, Sylvester J, Wilson K. American Medical Society of Sports Medicine Position Statement: Mononucleosis and Athletic Participation. Clin J Sport Med. 2023 May 15; [PubMed]

[8]. González-Andrade B, Santos-Lartigue R, Flores-Treviño S, Ramirez-Ochoa NS, Bocanegra-Ibarias P, Huerta-Torres FJ, Mendoza-Olazarán S, Villarreal-Treviño L, Camacho-Ortiz A, Villarreal-Vázquez H, Garza-González E. The carriage of interleukin-1B-31*C allele plus Staphylococcus aureus and Haemophilus influenzae increases the risk of recurrent tonsillitis in a Mexican population. PLoS One. 2017;12(5):e0178115. [PMC free article] [PubMed]

[9]. Alves METF, Mello LF, Alves FRF. Acute Epstein-Barr Virus and SARS-CoV-2 Coinfection: A Case Report. Curr Health Sci J. 2024 Jul-Sep;50(3):453-457. [PMC free article] [PubMed]

[10]. Miller JM, Binnicker MJ, Campbell S, Carroll KC, Chapin KC, Gonzalez MD, Harrington A, Jerris RC, Kehl SC, Leal SM, Patel R, Pritt BS, Richter SS, Robinson-Dunn B, Snyder JW, Telford S, Theel ES, Thomson RB, Weinstein MP, Yao JD. Guide to Utilization of the Microbiology Laboratory for Diagnosis of Infectious Diseases: 2024 Update by the Infectious Diseases Society of America (IDSA) and the American Society for Microbiology (ASM). Clin Infect Dis. 2024 Mar 05; [PubMed]

[11]. Pallon J, Rööst M, Sundqvist M, Hedin K. The aetiology of pharyngotonsillitis in primary health care: a prospective observational study. BMC Infect Dis. 2021 Sep 17;21(1):971. [PMC free article] [PubMed]

[12]. Andersen C, Greve T, Reinholdt KB, Kjaerulff AMG, Udholm N, Khalid V, Madzak A, Duez C, Münch H, Pauli S, Danstrup CS, Petersen NK, Klug TE. Bacterial findings in patients referred to hospital for the treatment of acute

tonsillitis with or without peritonsillar phlegmon. BMC Infect Dis. 2023 Jun 29;23(1):439. [PMC free article] [PubMed]

[13]. Wang Q, Du J, Jie C, Ouyang H, Luo R, Li W. Bacteriology and antibiotic sensitivity of tonsillar diseases in Chinese children. Eur Arch Otorhinolaryngol. 2017 Aug;274(8):3153-3159. [PubMed]

[14]. Ben Ammar C, Tbini M, Kamel HE, Riahi I, Ben Salah M. Tonsillar tuberculosis mimicking tonsillar carcinoma: A case report. Int J Surg Case Rep. 2024 Aug;121:110032. [PMC free article] [PubMed]

[15]. Klagisa R, Racenis K, Broks R, Balode AO, Kise L, Kroica J. Analysis of Microorganism Colonization, Biofilm Production, and Antibacterial Susceptibility in Recurrent Tonsillitis and Peritonsillar Abscess Patients. Int J Mol Sci. 2022 Sep 07;23(18) [PMC free article] [PubMed]

[16]. Banigo A, Moinie A, Bleach N, Chand M, Chalker V, Lamagni T. Have reducing tonsillectomy rates in England led to increasing incidence of invasive Group A Streptococcus infections in children? Clin Otolaryngol. 2018 Jun;43(3):912-919. [PubMed]

[17]. Douglas CM, Lang K, Whitmer WM, Wilson JA, Mackenzie K. The effect of tonsillectomy on the morbidity from recurrent tonsillitis. Clin Otolaryngol. 2017 Dec;42(6):1206-1210. [PMC free article] [PubMed]

[18]. Seidel DU, Bode S, Kostev K. Seasonal variation and demographic characteristics of acute tonsillitis and peritonsillar abscess in ENT practices in Germany (2019-2023). Eur Arch Otorhinolaryngol. 2025 Jan;282(1):317-326. [PubMed]

[19]. Stelter K. Tonsillitis and sore throat in children. GMS Curr Top Otorhinolaryngol Head Neck Surg. 2014;13:Doc07.

[20]. Sadeghirad B, Siemieniuk RAC, Brignardello-Petersen R, et al. Corticosteroids for treatment of sore throat: systematic review and metaanalysis of randomised trials. BMJ. 2017;358:j3887.

[21]. Brinckmann J, Sigwart H, van Houten Taylor L. Safety and efficacy of a traditional herbal medicine (Throat Coat) in symptomatic temporary relief of pain in patients with acute pharyngitis: a multicenter, prospective, randomized, double-blinded, placebo-controlled study. J Altern Complement Med. 2003;9(2):285-298.

[22]. Grimm W, Müller HH. A randomized controlled trial of the effect of fluid extract of Echinacea purpurea on the incidence and severity of colds and respiratory infections. Am J Med. 1999;106(2):138-143.

[23]. National Institute for Health and Care Excellence. Sore throat (acute): antimicrobial prescribing. January 2018. Accessed June 15, 2021. https://www.nice.org.uk/guidance/ng84/resources/visual-summarypdf-4723226606

[24]. van Driel ML, De Sutter AI, Thorning S, et al. Different antibiotic treatments for group A streptococcal pharyngitis. Cochrane Database Syst Rev. 2021;(3):CD004406.

[25]. Spinks A, Glasziou PP, Del Mar CB. Antibiotics for sore throat. Cochrane Database Syst Rev. 2013(11):CD000023.

[26]. Lexicomp. Accessed June 16, 2022. https://online.lexi.com/lco/action/ login

[27]. Trubiano JA, Vogrin S, Chua KYL, et al. Development and validation of a penicillin allergy clinical decision rule. JAMA Intern Med. 2020;180(5): 745-752.

[28]. Casey JR, Pichichero ME. Meta-analysis of cephalosporin versus penicillin treatment of group A streptococcal tonsillopharyngitis in children. Pediatrics. 2004;113(4):866-882.

[29]. Kocher JJ, Selby TD. Antibiotics for sore throat. Am Fam Physician. 2014;90(1):23-24.

[30]. Morad A, Sathe NA, Francis DO, McPheeters ML, Chinnadurai S. Tonsillectomy Versus Watchful Waiting for Recurrent Throat Infection: A Systematic Review. Pediatrics. 2017 Feb;139(2) [PMC free article] [PubMed]

[31]. Munck H, Jørgensen AW, Klug TE. Antibiotics for recurrent acute pharyngo-tonsillitis: systematic review. Eur J Clin Microbiol Infect Dis. 2018 Jul;37(7):1221-1230. [PubMed]

[32]. El Hennawi DED, Geneid A, Zaher S, Ahmed MR. Management of recurrent

tonsillitis in children. Am J Otolaryngol. 2017 Jul-Aug;38(4):371-374. [PubMed]

[33]. Wilson JA, O'Hara J, Fouweather T, Homer T, Stocken DD, Vale L, Haighton C, Rousseau N, Wilson R, McSweeney L, Wilkes S, Morrison J, MacKenzie K, Ah-See K, Carrie S, Hopkins C, Howe N, Hussain M, Mehanna H, Raine C, Sullivan F, von Wilamowitz-Moellendorff A, Teare MD. Conservative management versus tonsillectomy in adults with recurrent acute tonsillitis in the UK (NATTINA): a multicentre, open-label, randomised controlled trial. Lancet. 2023 Jun 17;401(10393):2051-2059. [PubMed]